

Report 609021064/2008

A.G. Oomen | P.J.C.M. Janssen | A. Dusseldorp | C.W. Noorlander

Exposure to chemicals via house dust



RIVM Report 609021064/2008

## Exposure to chemicals via house dust

A. G. Oomen P. J.C.M. Janssen A. Dusseldorp C. W. Noorlander

Contact: A. G. Oomen

Centre for Substances and Integrated Risk Assessment

E-mail: Agnes.Oomen@rivm.nl

This investigation has been performed by order and for the account of Inspectorate of the Ministry of Housing, Spatial Planning and the Environment of the Netherlands, within the framework of project M/609021.

# © RIVM 2008 Parts of this publication may be reproduced, provided acknowledgement is given to the 'National Institute for Public Health and the Environment', along with the title and year of publication.

RIVM Report 609021064

2

### **Abstract**

#### Exposure to chemicals via house dust

Humans are exposed to substances present in house dust, but the majority of these substances do not pose a risk to human health. Those substances that do exceed a critical level, however, have the potential to be a health risk. The most common of these are lead and di(2-ethylhexyl)phthalate, followed by arsenic, cadmium, polycyclic aromatic hydrocarbons and PBDEs (flame retardants). These substances come into house dust through the wear and tear of consumer products, the release of products when cooking and burning wood in the fireplace. In addition, some pollutants can fall off the soles of shoes that have come into contact with polluted soil.

By order of the Inspectorate of the Ministry of Housing, Spatial Planning and the Environment of the Netherlands, the Dutch National Institute for Public Health and the Environment (RIVM) has carried out a screening of the potential risks posed by various chemical substances in house dust. The substances investigated include metals, organotin compounds, phthalates, brominated flame retardants, pesticides and polycyclic aromatic hydrocarbons. Substances in house dust are mostly ingested through the contact of the hand or an object with the mouth. This is particularly true for young children. A minor part of house dust enters the body by inhalation.

The mean ingestion rates for adults and children were estimated. Exposure to substances via house dust is calculated based on the mean ingestion rates and typical concentrations of substances present in house dust. Whenever possible, this study has focussed on the situation in the Netherlands. The exposure of adults and children to substances via house dust was compared to the tolerable daily intake, which was used as a criterium for a potential health risk, and to the background exposure (via food and water consumption).

The findings of this investigation provide an overview of the substances in house dust which can exceed the accepted norm and for which the contribution of house dust to the total exposure is substantial. Based on these findings, the RIVM recommends that the substances identified herein be subjected to measurements in research on the indoor environment.

*Key words:* 

house dust; risk assessment; human health; children; exposure

## Rapport in het kort

#### Blootstelling aan chemische stoffen via huisstof

Mensen worden via huisstof aan chemische stoffen blootgesteld. De meeste stoffen vormen op deze manier geen risico voor de gezondheid. Voor enkele stoffen wordt wel de gezondheidskundige norm overschreden waardoor er mogelijk sprake kan zijn van een risico voor de gezondheid. Dit geldt met name voor lood en di(2-ethylhexyl)ftalaat en in minder mate voor arseen, cadmium, polycyclische aromatische koolwaterstoffen en PBDE's (vlamvertragers). Deze stoffen komen op allerlei manieren terecht in huisstof, bijvoorbeeld door slijtage van producten, inloop van verontreinigde bodem, door stoffen die bij het koken vrijkomen of via de open haard.

In opdracht van VROM-Inspectie heeft het RIVM een screening uitgevoerd van de risico's van verschillende chemische stoffen in huisstof (metalen, organotinverbindingen, ftalaten, gebromeerde vlamvertragers, bestrijdingsmiddelen, en polycyclische aromatische koolwaterstoffen). Huisstof wordt vooral ingenomen door contact van de hand of een voorwerp met de mond, wat vooral bij jonge kinderen veel voor komt. Daarnaast wordt een beperkte hoeveelheid huisstof ingeademd.

De inname van huisstof is geschat voor kinderen en volwassenen. De blootstelling aan chemische stoffen via huisstof is berekend op basis van de hoeveelheid huisstof die mensen binnenkrijgen en concentraties van chemische stoffen daarin. Waar mogelijk is dat op de situatie in Nederland toegespitst. De blootstelling via huisstof is vergeleken met de norm voor wat dagelijks is toegestaan, en met de achtergrondblootstelling via voeding en water.

De huidige bevindingen geven een overzicht van de stoffen in huisstof die de gezondheidskundige norm kunnen overschrijden, en waarvan de bijdrage van huisstof aan de totale blootstelling aanzienlijk is. Aanbevolen wordt de hier geïdentificeerde stoffen te meten bij onderzoek naar het binnenmilieu.

#### Trefwoorden:

Huisstof; risicobeoordeling; gezondheid; kinderen; blootstelling

## **Contents**

Sı	ımmary		7
1		Introduction	9
	1.1	Approach	9
	1.2	Assumptions	10
2		General information on house dust	11
	2.1	Origin of house dust	11
	2.2	Sampling	11
	2.3	Resuspension	11
3		Exposure to house dust	13
	3.1	Ingestion of house dust	13
	3.2	Inhalation of house dust	14
	3.3	Defaults used for house dust ingestion and inhalation	15
4		Intake of substances via house dust	17
	4.1	Chemicals	17
	4.1.1	Metals	17
	4.1.2	Pesticides	17
	4.1.3	Organotins	17
	4.1.4	Phthalates	18
	4.1.5	Brominated flame retardants	18
	4.1.6	Polycyclic aromatic hydrocarbons	18
	4.2	Levels of substances in house dust	18
5		Substances in house dust with and without potential human health risk	25
	5.1	Comparison exposure via house dust, TDI, and background exposure	25
	5.2	Levels of substances in particulate matter in air	30
	5.3	Substances without potential human health risk	31
	5.4	Substances with potential human health risk	31
6		Discussion substances with potential human health risk	35
	6.1	Aluminium	35
	6.2	Arsenic	36
	6.3	Cadmium	37
	6.4	Chromium	39
	6.5	Copper	39
	6.6	Lead	40
	6.7	Manganese	42
	6.8	Nickel	42
	6.9	Zinc	43
	6.10	Di(2-ethylhexyl)phthalate (DEHP)	44
	6.11	Sum PAHs	45
	6.12	Brominated flame retardants	47
	6.12.1	Toxicological reference values	47
	6.12.2	Ban on several brominated flame retardants	47

6.12.3	Possibility of a potential health risk for BDE99	47				
6.12.4	Comparison intake via house dust and food	48				
6.12.5	Comparison of exposure BDE209 via house dust to other congeners	49				
6.12.6	Other brominated flame retardants	49				
6.12.7	Non-brominated flame retardants	50				
6.13	Measures that can decrease the exposure to substances in house dust	50				
7	Conclusions	53				
Acknowled	gement	59				
References		61				
Appendix 1	. References on substances in house dust	71				
Appendix 2 List of TDIs and background exposure estimates						

6

## **Summary**

Up till now, various projects on the indoor environment remain inconclusive with regard to the human health risk of substances in house dust. From experience it is known that some substances such as lead in house dust may pose a human health risk, but it is unknown whether and to which extent this is also the case for other substances. In order to better design future projects on the indoor environment, the present investigation examines whether exposure to a great number of substances present in house dust may lead to potential human health risks.

House dust mainly enters the body by ingestion, especially for young children. Mean ingestion rates of house dust of 50 and 100 mg were estimated for adults and children, respectively. Typical concentrations of substances in house dust were extracted from literature. The exposure of a substance via house dust by adults and children was compared to the tolerable daily intake (TDI) as a criterium above which health risks may occur. The present research was restricted to several main chemical groups: metals, organotin compounds, pesticides, phthalates, brominated flame retardants and polycyclic aromatic hydrocarbons (PAHs). Based on calculation of the so-called Risk Index, which is the estimated exposure for any substance divided by its TDI, a list of substances is provided for which there may be a health risk.

We found the majority of substances present in house dust to be without potential human health risk. However, some substances in house dust exceed the TDI criterium, indicating that they may cause a potential risk. Substances which exceed this criterium most frequently are lead and di(2-ethylhexyl)phthalate, followed by arsenic, cadmium, sum PAHs, and BDE99. The finding that several metals (lead, arsenic, cadmium) may cause a potential human health risk via house dust was expected, but for the other substances this is not generally known. We recommend including these substances in the measurements when projects on the indoor environment are designed.

## 1 Introduction

House dust can be a major exposure route for some substances, leading to potential health risks. Especially young children ingest considerable amounts of house dust via hand-to-mouth and object-to-mouth behaviour. The ingestion of house dust by children is particularly high relative to their lower body weight. In addition, humans inhale dust particles with air, which may also contribute to the exposure.

The human health risks of chemical substances in house dust have only occasionally been investigated in the Netherlands. The present investigation examines whether exposure to a great number of substances present in house dust may potentially lead to human health risks. The research was restricted to several main chemical groups: metals, pesticides, including organotin compounds, phthalates, brominated flame retardants and polycyclic aromatic hydrocarbons. The aim of this research is to provide a list of substances that may cause human health risks due to exposure to house dust. Also substances are identified for which it is highly unlikely that they cause a human health risk via house dust. In assessing the possibility of health risks due to substances in soil, conditions that are representative for the Netherlands are pursued.

It should be noted that the present report does not assess the health risks associated with particulate matter (in Dutch 'fijn stof') in the air. Potential health risks associated with different *chemicals* in house dust and particulate matter are investigated.

Finally, some measures that can decrease the exposure to substances in house dust are discussed.

## 1.1 Approach

To investigate the potential health risk of substances in house dust, we used the following approach:

- o The rate of ingested and inhaled house dust by children and adults is estimated.
- o <u>Typical concentrations</u> of the substances in house dust are obtained from the literature wherever the research was performed.
- The toxicological reference levels (expressed as Tolerable Daily Intake) of the substances are identified as a criterium above which there may be a potential health risk. The TDI is expressed in mg per kg body weight per day. As contained within the definition of the TDI, lifelong exposure to the substance at levels below the TDI will in principle not lead to health effects. Exceptions and other approaches are indicated.
- o The <u>background exposures</u>, usually due to food and water consumption, of the substances are also listed.
- o The substances are identified for which it is highly unlikely that exposure via house dust can result in a possible health risk. This is accomplished by making a

conservative estimate of the human exposure (both adult and child) to the substance by using the highest encountered house dust concentration found in literature. Background exposure is added to the exposure via house dust and compared to the TDI, so that exposure via other routes (e.g. food) is included. When the added exposure via house dust and background is less than the TDI, a potential health risk is considered to be highly unlikely.

- The substances for which the added exposure via house dust (based on highest reported concentration) and background is higher than the TDI, are investigated in more detail to identify whether exposure via house dust indeed poses a potential health risk. This research includes:
  - estimation of the exposure with a 95<sup>th</sup>-percentile and a geometric mean concentration value in house dust
  - evaluation of the house dust data used in the calculations (can the data be considered to be representative for the Netherlands, are there enough data)
  - comparison of the exposure via house dust relative to other exposure routes (e.g. background exposure)
  - evaluation of the application of the TDI. In principle it is scientifically justified to integrate childhood and adult exposure. However, there are exceptions depending on the toxicological basis for the TDI.

In this manner the substances are identified for which exposure via house dust may lead to a potential health risk.

## 1.2 Assumptions

In order to assess the potential human health risk of substances in house dust, several assumptions have been made. These assumptions are:

- O Using the TDI as a toxic measure for comparison to the exposure to a substance, implies the assumption that humans are chronically exposed to that substance. It is realistic to assume that any person is daily exposed to a substance in house dust. However, changes in the concentration of the substance in house dust can occur in time. These changes are unknown and can therefore not be included in the calculation of the exposure.
- o The aim of the present investigation is to provide a *screening* whether exposure to house dust may lead to potential human health effects. With this aim in mind the estimation of the exposure to substances in house dust should be conservative.
- Where the present screening indicates a human health risk is unlikely, this does not mean that this cannot occur under any circumstances. Obviously, specific situations may still result in excessive concentrations in house dust of a certain substance leading to a health risk.

## 2 General information on house dust

#### 2.1 Origin of house dust

House dust is a heterogeneous mixture of substances from numerous sources, including tracked-in or resuspended soil particles, clothing, atmospheric deposition of particulates, hair, fibres (artificial and natural), molds, pollen, allergens, bacteria, viruses, arthropods, ash, soot, animal fur and dander, smoke, skin particles, cooking and heating residues, and building components among others ((Paustenbach *et al.*, 1997), and references herein).

## 2.2 Sampling

Reproducible sampling of house dust is difficult and highly depending on the method. Methods that are regularly used for dust collection are vacuum cleaners and wipes. Some studies suggest that the sampling efficiency can differ to a factor sometimes higher than 100 between methods (Sterling *et al.*, 1999). Wiping as sampling method is probably more reproducible, especially when wiping is only applied on hard surfaces. The effect of the sampling method on the concentration of the substance in house dust is unknown and probably depending on various factors.

## 2.3 Resuspension

Meyer *et al.* (1999) determined that the number of persons living in a residence was significantly associated with elevated amounts of dust sedimented per day, probably due to increased indoor activities such as vacuuming, sweeping, cleaning, and children playing. Thatcher and Layton have shown that the resuspension rate is particle size dependent (Thatcher and Layton, 1995). Particles with diameters of 5-25  $\mu$ m are most readily resuspended and even light activity such as walking into and out of the room can have a significant impact on the concentration of airborne particles greater than 5  $\mu$ m. Particles of 0.3 - 1  $\mu$ m, however, are not affected by either cleaning or walking (Thatcher and Layton, 1995).

## 3 Exposure to house dust

## 3.1 Ingestion of house dust

Several institutes have estimated the amount of house dust that children and adults ingest. These dust ingestion rates are uncertain as only indirect methods exist to estimate daily dust ingestion. Table 1 provides an overview of the estimated amounts and some information on the derivation of these estimations.

Table 1. House dust ingestion rates for adults and children derived and used in references.

Study	Ingestion house	e dust (mg/day)	Remarks		
	Adult	Children			
Gevao <i>et al.</i> , 2006	10	100	Based on Chuang et al., 1999		
Chuang <i>et al.</i> , 1999	60	100	Based on Lewis <i>et al.</i> , 1994; Stanek and Calabrese, 1995		
Williams, 2002	50	50-100	Exposure parameters, no derivation mentioned		
Calabrese et al., 1989	10-100	20-200	Based on soil ingestion		
Lewis <i>et al.</i> , 2001	2-10	20-100	Estimation based on inhalation and hand-mouth behaviour		
Butte et al., 2002	2-10	20-100	Estimation based on other studies		
USEPA, 1997	50	100	Based on soil ingestion		
Jones-Otazo <i>et al.</i> , 2005	20	50	From Health Canada 1994		
Maertens et al., 2004	0.56	50-100	Based on Hawley, 1985		
Yamamoto 2006	100	200	Based on Ministry of Environment, Japan 2001		
Wilford et al., 2005	4.16	55	Based on USEPA, 1997		
Stapleton et al., 2005		20-200	Based on Roberts and Dickey, 1995; USEPA, 2002		
Mushak, 1998		100-200	Intake based on other studies		
Oomen et al., 2007	39	27	Based on a comparison with soil ingestion rates and hand loading*		
Range of studies	0.56-100	20-200			

<sup>\*</sup> Assuming children to remain outdoor during 2.9 h/day, and adults during 1.1 h/day.

In the present study a conservative but realistic estimate of dust ingestion rate will be used for the calculation of human exposure to substances in dust. This will allow identification of those substances for which it is unlikely that exposure via house dust will cause a human health risk.

As a conservative but realistic estimate of dust ingestion 100 mg/day for children and 50 mg/day for adults will be employed. This expert judgement is based on two arguments. Firstly, it is assumed in the Netherlands that children ingest on average 100 mg *soil* per day via hand-to-mouth behaviour (Lijzen *et al.*, 2001; Otte *et al.*, 2001). When playing outside a child's hand is much more loaded with soil than a loading with house dust during indoor playing. The ingestion of soil per time unit will thus be much greater outdoors than indoors. On the other hand, children spend more time indoors than outdoors. Yet, it is very unlikely that average daily dust ingestion will be greater than average daily soil ingestion.

Secondly, Table 1 shows that in some cases the dust ingestion of children is estimated at 200 mg/day, whereas in most cases 100 mg/day is used as upper level. For adults, in most cases about 50 mg/day was derived as an upper estimate.

#### 3.2 Inhalation of house dust

The amount of inhaled house dust can be estimated from the level of particles in the air  $(mg/m^3)$  and the volume of air inhaled by a child or adult. It is generally assumed that a child inhales 7.6 m<sup>3</sup> of air daily and an adult 19.9 m<sup>3</sup> (Otte *et al.*, 2001; Lewis *et al.*, 1999). As default values for body weight 15 kg and 70 kg respectively are used (Otte *et al.*, 2001).

In general, particles with an aerodynamic diameter of 10  $\mu$ m or smaller can be inhaled by humans and can deposit deep in the lungs (Carrizales *et al.*, 2005). Larger particles mainly deposit in the upper bronchial tubes and in most cases are transferred upwards by mucotransilliarly and subsequently ingested. Therefore, it is assumed that all substances associated with particles of 10  $\mu$ m in size or smaller contribute to the exposure in the lungs. This may be an overestimation as only part of the particles will be deposited in the lungs.

Concentrations of suspended particles in air, i.e. suspended dust, that are typically observed range between 13 and 35  $\mu$ g/m³ inside homes, between 41 and 58  $\mu$ g/m³ for daycare facilities (Beamer *et al.*, 2002). Higher concentrations are usually found directly around persons (personal cloud) than in other places in a room. A value of 60  $\mu$ g/m³ is probably representative for moderately crowded places such as residents, whereas a value of 100  $\mu$ g/m³ should be used as personal exposure for crowded places such as classrooms (Oomen and Lijzen., 2004).

When assuming a constant concentration of suspended particles in air of  $100~\mu g/m^3$ , and a volume of inhaled air of  $7.6~m^3$  for a child and  $19.9~m^3$  for an adult, the amount of inhaled suspended particles are respectively 0.8 and 2.0~mg per day. This is in line with the inhaled amounts of dust reported by Maertens *et al.* (2004). Hence, the amount of inhaled suspended dust particles is low compared to the amount of ingested dust (50 and 100 mg/day for a child and adult, respectively). When considering the total exposure to a substance in house dust exposure via inhalation is negligible. Obviously, this exposure route should still be considered when a substance may have local effects on the lungs. For example for several metals this may be the case. In order to assess whether concentrations of metals in the indoor air may lead to potential health risks, concentrations in airborne particulate matter were retrieved from literature and compared to Tolerable Concentration in Air (TCA) levels  $(ng/m^3)$ , see section 5.2. It should be stressed that the literature research on indoor air concentrations for metals was limited. At concentrations of airborne particulate matter below the TCA no health risk is anticipated even after life-long exposure.

Furthermore, it should be noted that substances that enter the blood circulation after deposition in the lungs bypass the first pass effect, i.e. these substances directly enter the systemic blood circulation without possible metabolism in the liver as is the case after oral exposure. This difference may have consequences for the toxicity.

Obviously, volatile substances are not necessarily adhered to dust particles so that information on the concentration of the substance in airborne particulate matter is not useful.

## 3.3 Defaults used for house dust ingestion and inhalation

Based on section 3.1 and 3.2 the default dust ingestion and inhalation values described in Table 2 are used in the calculation of the human exposure to substances in house dust. Due to the large uncertainties in these default values, and the small contribution of inhalation to the total house dust intake, a total daily intake of house dust of 50 and 100 mg is used for adults and children, respectively.

Table 2. Default values for house dust ingestion and inhalation

	Ingestion of house dust	Inhalation of house dust	Total intake of house
	(mg/day)	(mg/day)	dust (mg/day)
Adult	50	0.8	50
Child	100	2.0	100

## 4 Intake of substances via house dust

#### 4.1 Chemicals

Many chemicals are present in house dust as a consequence of their widespread use in everyday consumer products present in homes. The present research is restricted to the analysis of five main compound groups covering the chemicals: metals, organotins, pesticides (including organotins), phthalates, brominated flame retardants and polycyclic aromatic hydrocarbons.

#### **4.1.1 Metals**

Metals present at trace levels in natural water, air, dusts, soils and sediments, play an important role in human life (Juvanovic *et al.*, 1995; Lapitajs *et al.*, 1995). Sources of trace elements in house dust are atmospheric fall out of petrol, tyre wear, corrosion of metallic parts of automobiles, rooftiles, paint and release from carpets, smoking (Fergusson and Schroeder, 1985; Fergusson and Kim, 1991).

#### 4.1.2 Pesticides

Pesticides may be a chemical substance, biological agent, antimicrobial, disinfectant or device used against any pest. There are several types of pesticides, such as bactericides, fungicides, insecticides and herbicides.

#### 4.1.3 Organotins

Organotins are chemical compounds based on tin with hydrocarbon substituents. Organotins are primarily used in five major commercial applications: PVC heat stabilizers, biocides, catalysts, agrichemicals and glass coatings. Triorganotins have a high toxicity and can be powerful fungicides and bactericides, depending on the organic group present. Tributyltins are industrial biocides used in antifouling paints and in wood treatment and preservation. Tributyltins are also used as disinfectants, molluscicides, antifungal action in textiles and industrial water systems such as cooling tower and refrigeration water systems, wood pulp and paper mill systems, and breweries. Many of these applications have been phased out because of the high aquatic toxicity of tributyltins. Triphenyltins are used as fungicides, miticides and acaricides.

#### 4.1.4 Phthalates

Phthalates, or phthalate esters, are a group of chemicals that are mainly used as softeners in flexible PVC products. Phthalates can be found in a broad range of consumer products like packaging materials, wallpapers, furnishings, clothing and toys, as well as ingredients in cosmetics and perfumes.

#### 4.1.5 Brominated flame retardants

Brominated flame retardants (BFRs) are a group of brominated organic substances that have an inhibitory effect on the ignition of combustible organic materials. The most widely used BFRs are tetrabromobisphenol-A (TBBP-A), hexabromocyclododecane (HBCDD), and polybrominated diphenylethers (PBDEs) (De Winter-Sorkina *et al.*, 2006). BFRs are applied to textiles, wiring, furniture, industrial paints and incorporated into plastics and foams, and they are commonly used in electronic products to reduce the flammability of the product.

Use of pentabromodiphenylether (penta-BDE) technical product was voluntarily phased out by industry within the European Union over the last 10 years. This has led to increased use of HBCDD and TBBP-A. The use of penta-BDE and octa-BDE technical products in all applications for the European Union market has officially been banned since August 2004. The use of penta-BDE, octa-BDE and PBBs in new electrical and electronical equipment is banned from July 2006 (De Winter-Sorkina *et al.*, 2006).

#### 4.1.6 Polycyclic aromatic hydrocarbons

Polycyclic aromatic hydrocarbons (PAHs) are a group of chemicals composed of two or more fused aromatic rings made up of carbon and hydrogen. Polycyclic aromatic hydrocarbons are formed by the incomplete combustion of coal, oil, petrol, wood, tobacco, charbroiled meats, garbage, or other organic materials. A few are used in medicines, and to make dyes, plastics, and pesticides. Naphthalene, is used in making dyes, explosives, plastics, lubricants, and moth repellent. Anthracene is used in dyes, insecticides and wood preservatives. PAHs are present in tobacco smoke, smoke from home heating (burning wood or oil), char-grilled food and creosote treated wood products. High concentrations of polycyclic aromatic hydrocarbons have been found in coal-tar production plants, coking plants, bitumen and asphalt production plants, smoke houses, aluminium production plants, and trash incinerators. PAHs are also present in the soil where coal, wood, petrol or other products have been burned. Food produced from these soils may also contain PAHs. Most of the PAHs are considered genotoxic carcinogens.

#### 4.2 Levels of substances in house dust

An overview of concentrations of chemicals in house dust as obtained from literature is presented in Table 3. The highest concentration found in literature is presented. In

addition, for most substances several geometric mean values are found in literature (from different studies). In the present report the highest geometric means (mg/kg) of the concentrations of chemicals in house dust under normal conditions are used. Using these concentration values, ingestion of chemicals (µg/kg body weight/day) was calculated for both adults and children. For this calculation mean body weights of 70 kg and 15 kg were used for adults and children respectively (Otte *et al.*, 2001) and dust ingestion rates of 50 mg/day for adults and 100 mg/day for children were used.

Table 3. Calculated ingestion rates of substances via house dust for an adult (70 kg) and child (15 kg) based on the highest geometric mean and the maximum concentration of the compound in house dust described in literature (see Appendix 1 for additional information and references).

	Geometric mean	Mean ingestion rate		Maximum	Maximum ingestion rate		
Compound		Adult	Child		Adult	Child	
	(mg/kg)	(μg/kg/d)	(µg/kg/d)	(mg/kg)	(µg/kg/d)	(µg/kg/d)	
Metals		•	•		•	•	
Aluminium	24281	17	162	51100	36	341	
Antimony	26	0.018	0.17	66	0.05	0.44	
Arsenic	70	0.050	0.47	192	0.14	1.3	
Barium	454	0.32	3.0	1480	1.1	9.9	
Beryllium	0.53	0.0004	0.0035	1	0.001	0.01	
Bismuth	1.0	0.0007	0.0068	8.6	0.01	0.06	
Cadmium	13	0.009	0.087	220	0.16	1.5	
Chromium <sup>1</sup>	159	0.11	1.1	5440	3.9	36	
Cobalt	17	0.012	0.11	23	0.02	0.15	
Copper	261	0.19	1.7	12540	8.9	84	
Lead	1200	0.85	8.0	37000	26	247	
Lithium	6.1	0.0043	0.041	16	0.01	0.10	
Magnesium	9442	6.7	63	52000	37	347	
Manganese	260	0.18	1.7	9410	6.7	63	
Mercury	1.7	0.0012	0.012	37	0.03	0.25	
Molybdenum	2.8	0.0020	0.019	29	0.02	0.19	
Nickel	47	0.033	0.31	243	0.17	1.6	
Rubidium	25	0.017	0.16	40	0.03	0.27	
Selenium	1.0	0.0007	0.007	6.8	0.005	0.05	
Silver	1.5	0.0011	0.010	9.3	0.01	0.06	
Strontium	242	0.17	1.6	1170	0.83	7.8	
Tellurium	0.07	0.00005	0.0005	0.28	0.0002	0.002	
Thallium	0.14	0.0001	0.0009	0.24	0.0002	0.002	
Tin	22	0.016	0.15	595	0.42	4.0	

	Geometric mean	Mean inge	estion rate	Maximum	Maximum ingestion rate		
Compound		Adult	Child		Adult	Child	
	(mg/kg)	(µg/kg/d)	(μg/kg/d)	(mg/kg)	(μg/kg/d)	(μg/kg/d)	
Metals			•	1	ı		
Titanium	2854	2.0	19	4983	3.5	33	
Tungsten	3.7	0.003	0.025	5.6	0.004	0.04	
Uranium	0.55	0.0004	0.004	1.3	0.001	0.01	
Vanadium	112	0.080	0.75	193	0.14	1.3	
Zinc <sup>2</sup>	628	0.45	4.2	30600	22	204	
Organotin compour	nds						
Dibutyltin (DBT)	0.51	0.0004	0.0034	5.6	0.004	0.04	
Dioctyltin (DOT)	0.02	0.00001	0.0001	0.36	0.0003	0.002	
Monobutyltin (MBT)	0.16	0.0001	0.0011	1.5	0.001	0.01	
Monooctyltin (MOT)	0.01	0.00001	0.0001	0.04	0.00003	0.0003	
Tributyltin (TBT)	0.02	0.00001	0.0001	0.08	0.0001	0.001	
Sum organotins <sup>3</sup>	0.7	0.0005	0.005	7.2	0.005	0.05	
Pesticides							
2,4- Dichlorophenoxy acetic acid	1.24	0.0009	0.0083	7.3	0.005	0.05	
Alachlor <sup>4</sup>				1.5	0.001	0.01	
Aldrin	0.006	0.000004	0.00004	0.051	0.00004	0.0003	
alpha-Chlordane <sup>5</sup>	0.055	0.00004	0.0004	0.26	0.0002	0.002	
alpha-HCH	0.0007	0.000000	0.000005	0.0087	0.00001	0.0001	
Atrazine	0.002	0.000002	0.00002				
Azinphos methyl	6.0	0.0043	0.040	16	0.01	0.11	
beta-HCH	0.0022	0.000002	0.00001	0.057	0.00004	0.0004	
Carbaryl <sup>4</sup>				1.0	0.0007	0.007	
Chloroprofam <sup>4</sup>				0.17	0.0001	0.001	
Chlorpyrifos	1.0	0.0007	0.0069	6.5	0.005	0.04	
DDD	0.0047	0.000003	0.00003	0.048	0.00003	0.0003	
DDE	0.007	0.000005	0.00005	0.05	0.00004	0.0003	
DDT	0.12	0.0001	0.0008	0.78	0.0006	0.005	
delta-HCH	0.0055	0.000004	0.00004	0.17	0.0001	0.001	
Diazinon	0.31	0.0002	0.0021	2.0	0.001	0.01	
Dicamba <sup>4</sup>				2.5	0.002	0.02	

	Geometric mean	Mean inge	estion rate	Maximum	Maximum ingestion rate		
Compound		Adult	Child		Adult	Child	
-	(mg/kg)	(μg/kg/d)	(μg/kg/d)	(mg/kg)	(μg/kg/d)	(μg/kg/d)	
Pesticides	, , ,	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	1, 5 5	,, ,	1.00	
Dieldrin	0.018	0.00001	0.0001	0.050	0.00004	0.0003	
Dimethyl	0.37	0.0003	0.0025	1.3	0.0009	0.009	
organiphosphate							
Ethyl parathion	0.56	0.0004	0.0037	0.43	0.0003	0.003	
gamma-Chlordane <sup>6</sup>	0.098	0.0001	0.0007	0.47	0.0003	0.003	
Glyphosate	0.14	0.0001	0.0009	0.14	0.0001	0.001	
Heptachlor	0.12	0.0001	0.0008	0.34	0.0002	0.002	
Lindane <sup>7</sup>	0.33	0.0002	0.0022	0.074	0.0001	0.0005	
Malathion	0.38	0.0003	0.0025	2.0	0.001	0.013	
Mecoprop <sup>4</sup>				0.40	0.0003	0.003	
Methamidophos <sup>4</sup>				0.40	0.0003	0.003	
Methyl parathion	0.38	0.0003	0.0025	1.9	0.0013	0.013	
Metolachlor	0.0057	0.000004	0.00004	0.80	0.0006	0.005	
Pendimethalin <sup>4</sup>	0.0007	0.00000	0.0000.	3.0	0.002	0.020	
Permethrin	0.14	0.0001	0.0009	659	0.47	4.4	
Phosmet	5.2	0.0037	0.035	22	0.016	0.15	
Picloram <sup>4</sup>	3.2	0.0037	0.033	1.2	0.0009	0.008	
Resmethrin <sup>4</sup>				0.80	0.0006	0.005	
Tetramethrin <sup>4</sup>				0.80	0.0003	0.003	
Trichloro-2-	0.54	0.0004	0.0036	0.40	0.0003	0.003	
pyridinol <sup>8</sup>	0.54	0.0004	0.0036	0.93	0.0007	0.006	
Trifluralin <sup>4</sup>				1.8	0.0013	0.012	
Phthalates			1		1		
Butylbenzyl	319	0.23	2.1	45549	32	304	
phthalate (BBP)					_	= * *	
Di(2-ethylhexyl)	3214	2.3	21	40459	29	270	
phthalate (DEHP)						1	
Diethyl phthalate	45	0.032	0.30	632	0.45	4.2	
(DEP) Diisobutyl phthalate	84	0.060	0.56	84	0.06	0.56	
(DiBP)	07	0.000	0.30	04	0.00	0.50	
Diisodecyl phthalate	73	0.052	0.49	73	0.05	0.49	
(DIDP)							
Diisononyl	176	0.12	1.2	176	0.12	1.2	
phthalate (DINP)	11	0.00==	0.072	1.50	0.11	1.1	
Dimethyl phthalate (DMP)	11	0.0077	0.072	158	0.11	1.1	

	Geometric mean	Mean ingestion rate		Maximum	Maximum rate	ingestion
Compound		Adult	Child		Adult	Child
	(mg/kg)	(μg/kg/d)	(μg/kg/d)	(mg/kg)	(μg/kg/d)	(µg/kg/d)
Phthalates			1	1	•	<u> </u>
Dimethylpropyl phthalate (DMPP)	55	0.039	0.36	161	0.11	1.1
Di-n-butyl phthalate (DBP)	226	0.16	1.5	5446	3.9	36
Brominated flame ret	tandants (DED	Ps)/Puomina	tad dinhamil	ath aus (PDEs)		
BDE 100	490	0.35	3.3	21000	15	140
BDE 138	37	0.33	0.25	2000	1.4	13
BDE 158	181	0.03	1.2	1510	1.4	10
				_		
BDE 154 BDE 17	380	0.27	2.5	18000 150	13	120
BDE 17 BDE 183	8.9	0.01	0.06	650	0.1	1.0
		0.03				4.3
BDE 190	4.5	0.003	0.03	48	0.0	0.3
BDE 196	15	0.01	0.10	39	0.0	0.3
BDE 197	17	0.01	0.12	77	0.1	0.5
BDE 206	51	0.04	0.34	239	0.2	1.6
BDE 207	30	0.02	0.20	109	0.1	0.7
BDE 208	35	0.02	0.23	108 19100	0.1	0.7
BDE 209	10	0.01				127
BDE 28	20	0.01	0.14	550	0.4	3.7
BDE 33/28	21	0.01	0.14	77	0.1	0.5
BDE 47	1621	1.2	11	33	0.0	0.2
BDE 66	37	0.03	0.25	1800	1.3	12
BDE 85	190	0.13	1.3	9700	7	65
BDE 99	2295	1.6	15	2850	2	19
BDE153	470	0.33	3.1	25000	18	167
Polycyclic aromatic h	ydrocarbons	(PAHs)				
Acenaphthene	0.05	0.00004	0.0003	1.9	0.001	0.013
Acenaphthylene	0.08	0.0001	0.0005	0.52	0.0004	0.003
Anthracene	0.12	0.0001	0.0008	5.8	0.004	0.039
Benz[a]anthracene	0.24	0.0002	0.0016	40	0.028	0.27
Benzo[a]pyrene	0.29	0.0002	0.0019	54	0.038	0.36
Benzo[b,k]fluoranth ene	0.57	0.0004	0.0038	108	0.077	0.72
Benzo[e]pyrene	0.29	0.0002	0.0019	41	0.029	0.27
Benzo[g,h,i]perylene	0.25	0.0002	0.0017	35	0.025	0.23

	Geometric mean	Mean ingestion rate		Maximum	Maximum ingestion rate	
Compound		Adult	Child		Adult	Child
	(mg/kg)	(µg/kg/d)	(μg/kg/d)	(mg/kg)	(μg/kg/d)	(µg/kg/d)
Polycyclic aromatic h	ydrocarbons (	(PAHs)		•		
Biphenyl	0.002	0.000001	0.00001	0.005	0.000004	0.00003
Chrysene	0.39	0.0003	0.0026	43	0.031	0.29
Coronene	0.13	0.0001	0.0009	7.2	0.005	0.048
Cyclopenta[c,d]pyrene	0.08	0.0001	0.0005	0.62	0.0004	0.004
Dibenzo[a,h]anthracene	0.10	0.0001	0.0007	9.0	0.006	0.060
Fluoranthene	0.59	0.0004	0.0039	90	0.064	0.60
Fluorene	0.12	0.0001	0.0008	3.0	0.002	0.020
Indeno[1,2,3-c,d]pyrene	0.26	0.0002	0.0017	41	0.029	0.27
Naphthalene	0.33	0.0002	0.0022	42	0.030	0.28
Phenanthrene	0.44	0.0003	0.0029	43	0.031	0.29
Pyrene	0.49	0.0003	0.0033	69	0.049	0.46
Sum PAHs <sup>9</sup>	4.8	0.0034	0.032	634	0.45	4.2

#### Table 3 – Remarks

- 1. From a toxicological point of view chromium should be differentiated into hexavalent chromium, soluble trivalent chromium and unsoluble chromium. No differentiation, however, was made in the papers, which have measured chromium (only total chromium levels measured).
- 2. Non industrial setting.
- 3. Sum of organotins: dibutyltin (DBT), monobutyltin (MBT), monooctyltin (MOT), tributyltin (TBT).
- 4. Of the compounds alachlor, carbaryl, chloroprofam, dicamba, mecoprop, methamidos, pendimethalin, picloram, resmethrin, tetramethrin and trifluralin only maximum values were found in literature.
- 5. Alpha-Chlordane = cis-Chlordane.
- 6. Gamma-Chlordane = trans-Chlordane.
- 7. Lindane is technical gamma-HCH.
- 8. Trichloro-2-pyridinol is a metabolite of chlorpyrifos.
- 9. Sum of PAHs: acenaphthene, acenaphthylene, anthracene, benzo[a]anthracene, benzo[a]pyrene, benzo[b,k]fluoranthene, benzo[e]pyrene, benzo[g,h,i]perylene, biphenyl, chrysene, coronene, cyclopenta[c,d]pyrene, dibenzo[a,h]anthracene, fluoranthene, fluorine, indeno[1,2,3-c,d]pyrene, naphthalene, phenanthrene, pyrene.

# 5 Substances in house dust with and without potential human health risk

# 5.1 Comparison exposure via house dust, TDI, and background exposure

In order to select the compounds for which it is highly unlikely that exposure via house dust results in a human health risk, the exposure via house dust was estimated in a conservative manner: by using the highest and the highest geometric mean concentration encountered in literature. As a toxicologically-based reference value the Tolerable Daily Intake (TDI) is used since this represents the estimated amount of the chemical that humans can ingest daily during their lifetime without resultant adverse effects. It is thus implicitly assumed that humans are exposed to house dust daily, and that the concentration of the compound in house dust does not change over time.

The TDI represents the upper limit of allowable exposure to a single compound. Exposure to this compound can occur via different exposure routes. Since ingestion of house dust occurs in addition to other exposure routes the background exposure, e.g. exposure via normal food and water consumption, has to be taken into account.

For convenience, a *Risk Index (RI)* is used to identify the possible health risk. The risk index (RI) is calculated by:

This means that background exposure is incorporated in the calculation of the potential health risk. For the RI, the mean and maximum ingestion rates were used for both adults and children (see Table 4). When RI < 1, no risk is expected for human health. However, when RI > 1 this indicates a potential risk for human health. Although it is unlikely that a substance can cause a health risk at RI < 1, an additional safety precaution was taken at this point to further investigate substances for which an RI > 0.8 was obtained. *Hence, all compounds with a risk index greater than 0.8 are addressed in more detail in section 5.3.* The main purpose of Table 4 is to identify those compounds for which it is highly unlikely that they can cause a potential health risk via exposure through house dust, even if a conservative approach is used (highest available concentration in house dust is used in the calculation of the human exposure, additional research for compounds with a RI > 0.8, accounting for background exposure).

The TDIs and background exposure used are listed in Appendix 2, including the references to their derivation.

Table 4. Tolerable daily intake and risk index for compounds present in house dust based on highest geometric mean and maximum levels in house dust described in literature (see also Table 3). Risk Indices (RI) greater than 0.8 were taken to indicate a potential health risk (bold values). The potential health risk for these compounds is discussed in more detail in section 5.3. Mean and maximum concentrations of substances in house dust used for the calculation can be found in Table 3.

Compound	TDI	Backg	round	Risk index				
	(µg/kg/d)	(μg/kg	/d)	mean adult	mean child	max adult	max child	
		Adult	Child					
Metals								
Aluminium <sup>1</sup>	750	$180^{2}$	300	0.26	0.62	0.29	0.85	
Antimony <sup>1</sup>	6.0	$0.48^{3}$	0.5	0.08	0.12	0.09	0.16	
Arsenic <sup>1</sup>	1.0	0.3	$0.7^{4}$	0.35	1.2	0.44	2.0	
Barium	600	9	.0	0.02	0.02	0.02	0.03	
Beryllium	0.5	0	.3	0.60	0.61	0.60	0.61	
Bismuth <sup>5</sup>	n.a. <sup>5</sup>		nown					
Cadmium	0.5	0.	45	0.92	1.1	1.2	3.8	
Chromium III soluble <sup>6</sup>	5.0	1	.0	0.22	0.41	0.97	7.5	
Chromium III insoluble <sup>6</sup>	5000	1	.0	0.00	0.00	0.00	0.01	
Chromium VI <sup>7</sup>	5.0	0	.0	0.00	0.00	0.00	0.00	
Cobalt	1.4	0	.6	0.44	0.51	0.44	0.54	
Copper	83	6	0	0.73	0.74	0.83	1.7	
Lead <sup>1</sup>	3.6	1.1	1.8	0.54	2.7	7.7	69	
Lithium <sup>5</sup>	n.a. <sup>5</sup>	unkr	nown					
Magnesium	6700	46	00	0.69	0.70	0.69	0.74	
Manganese	160	13	30	0.81	0.82	0.85	1.2	
Mercury	2.0	0	.1	0.05	0.06	0.06	0.17	
Molybdenum	10	4	.0	0.40	0.40	0.40	0.42	
Nickel <sup>1</sup>	10	4.0	8.0	0.40	0.83	0.42	0.96	
Rubidium <sup>5</sup>	n.a. <sup>5</sup>	unkr	nown					
Selenium	5.0	2	.0	0.40	0.40	0.40	0.41	
Silver	5.0	1.3		0.26	0.26	0.26	0.27	
Strontium	600	18		0.03	0.03	0.03	0.04	
Tellurium	2.0		4 <sup>14</sup>	0.70	0.70	0.70	0.70	
Thallium	0.2		13 <sup>14</sup>	0.15	0.15	0.15	0.16	
Tin	2000		90	0.15	0.15	0.15	0.15	
Titanium	12000	7	.0	0.001	0.002	0.001	0.003	

Compound	TDI	Background	Risk index				
	(µg/kg/d)	(µg/kg/d)	mean adult	mean child	max adult	max child	
		Adult Child					
Metals	<b>-</b>						
Tungsten <sup>5</sup>	n.a. <sup>5</sup>	unknown					
Uranium	2.0	0.06	0.03	0.03	0.03	0.03	
Vanadium	2.0	$0.3^{14}$	0.19	0.52	0.22	0.79	
Zinc <sup>8</sup>	500	350	0.70	0.71	0.74	1.1	
Organotin compounds							
Dibutyltin (DBT)							
Dioctyltin (DOT)							
Monobutyltin (MBT)							
Monooctyltin (MOT)							
Tributyltin (TBT)							
Sum organotins <sup>9</sup>	0.25	0.083	0.33	0.35	0.35	0.53	
Pesticides							
2,4-Dichlorophenoxy acetic acid	10	unknown	< 0.0001	0.0008	0.0005	0.0049	
Alachlor	10	unknown			0.0005	0.0049	
Aldrin	0.115	< 0.04	< 0.0001	0.0004	0.0004	0.0034	
alpha-Chlordane <sup>10</sup>	0.5	unknown	0.0001	0.0007	0.0004	0.0034	
alpha-HCH	1.0	< 0.03	< 0.0001	< 0.0001	< 0.0001	0.0001	
Atrazine	35	unknown	< 0.0001	< 0.0001			
Azinphos methyl	5	unknown	0.0009	0.0080	0.0023	0.0213	
beta-HCH	0.02	< 0.01	0.0001	0.0007	0.0020	0.0190	
Carbaryl	8	unknown			0.0001	0.0008	
Chloroprofam	50	unknown			< 0.0001	0.0000	
Chlorpyrifos	10	unknown	0.0001	0.0007	0.0005	0.0043	
DDD	0.5 <sup>16</sup>	unknown	< 0.0001	0.0001	0.0001	0.0006	
DDE	0.5 <sup>16</sup>	unknown	< 0.0001	0.0001	0.0001	0.0007	
DDT	0.5 <sup>16</sup>	unknown	0.0002	0.0016	0.0011	0.0104	
delta-HCH <sup>5</sup>	n.a. <sup>5</sup>	unknown					
Diazinon	5	unknown	< 0.0001	0.0004	0.0003	0.0027	
Dicamba	125	unknown			< 0.0001	0.0001	
Dieldrin	0.115	unknown	0.0001	0.0012	0.0004	0.0033	
Ethyl parathion	4	unknown	0.0001	0.0009	0.0001	0.0007	
gamma-Chlordane <sup>11</sup>	0.5	unknown	0.0001	0.0013	0.0007	0.0063	
Glyphosate	1000	unknown	< 0.0001	< 0.0001	< 0.0001	< 0.0001	

Compound	TDI	Background		Risk index				
	(µg/kg/d)	(μg/kg	/d)	mean adult	mean child	max adult	max child	
		Adult	Child					
Pesticides								
Heptachlor	0.1	0.0	001	0.0108	0.0179	0.0124	0.0323	
Lindane <sup>12</sup>	0.04			0.0059	0.0550	0.0013	0.0123	
Malathion	300	unkr	nown	< 0.0001	< 0.0001	< 0.0001	< 0.0001	
Mecoprop	3.3	unkr	nown			0.0001	0.0008	
Methamidophos	4	unkr	nown			0.0001	0.0007	
Methyl parathion	3	unkr	nown	0.0001	0.0008	0.0004	0.0042	
Metolachlor	3.5	unkr	nown	< 0.0001	< 0.0001	0.0002	0.0015	
Pendimethalin	125	unkr	nown			< 0.0001	0.0002	
Permethrin	50	unkr	nown	< 0.0001	< 0.0001	0.0094	0.0879	
Phosmet	3	unkr	nown	0.0012	0.0116	0.0052	0.0489	
Picloram	200	unkr	nown			< 0.0001	0.0000	
Resmethrin	30	unkr	nown			< 0.0001	0.0002	
Tetramethrin	20	unkr	nown			< 0.0001	0.0001	
Trichloro-2-pyridinol <sup>5</sup>	n.a. <sup>5</sup>	unkr	nown					
Trifluralin	15	unkr	nown			0.0001	0.0008	
Phthalates	l	I		l	l		l	
Butylbenzyl phthalate (BBP)	500	9.	00	0.02	0.02	0.08	0.63	
Di(2-ethylhexyl) phthalate (DEHP)	50	16 <sup>22</sup>	$26^{23}$	0.37	0.95	0.90	5.9	
Diethyl phthalate (DEP)	$200^{14}$	unkr	nown	0.0002	0.002	0.003	0.03	
Diisobutyl phthalate (DiBP) <sup>5</sup>	n.a. <sup>18</sup>	unkr	nown	see DBP	see DBP	see DBP	see DBP	
Diisodecyl phthalate (DIDP)	150	unkr	nown	0.0003	0.03	0.0003	0.00	
Diisononyl phthalate (DINP)	150	unkr	nown	0.001	0.01	0.001	0.01	
Dimethyl phthalate (DMP) <sup>5</sup>	n.a. <sup>17</sup>	unkr	nown	see DEP	see DEP	see DEP	see DEP	
Dimethylpropyl phthalate (DMPP) <sup>5</sup>	n.a. <sup>18</sup>	unkr	nown	see DBP	see DBP	see DBP	see DBP	
Di-n-butyl phthalate (DBP)	52	unknown		0.00	0.05	0.08	0.73	
( /								
Brominated flame retards	ants (BFRs)/	Bromin	ated dip	henylethers (E	BDEs)	1	1	
BDE 47 <sup>5</sup>	n.a. <sup>5</sup>	A: 0.0	$0030^{20} \\ 0014^{20}$	19	19	19	19	
BDE 99 - EU	0.00026	A: 0.0	$   \begin{array}{r}     \hline     0010^{20} \\     0023^{20}   \end{array} $	0.45	1.5	22	204	

Compound	TDI	Background	l Risk index				
	(µg/kg/d)	(µg/kg/d)	mean adult	mean child	max adult	max child	
		Adult Child					
Brominated flame retarda	ints (BFRs)	Brominated dip	henylethers (E	BDEs)			
BDE 99 - VS		A: 0.00010 <sup>20</sup>					
	0.00026	C: 0.00023 <sup>20</sup>	<b>6.8</b>	61	39	362	
BDE 100 <sup>5</sup>	n.a. <sup>5</sup>	A: 0.00007 <sup>20</sup>	19	19	19	19	
DDE 102 <sup>5</sup>	5	C: 0.00018 <sup>20</sup>	19	19	19	19	
BDE 183 <sup>5</sup>	n.a. <sup>5</sup>	A: 0.00034 <sup>20</sup> C: 0.00087 <sup>20</sup>					
BDE 209 <sup>5</sup>	n.a. <sup>5</sup>	C: 0.00087 <sup>20</sup>					
Polycyclic aromatic hydro		4Hs)			•		
Acenaphthene <sup>5</sup>	n.a. <sup>5</sup>						
Acenaphthylene <sup>5</sup>	n.a. <sup>5</sup>						
Anthracene	40						
Benz[a]anthracene <sup>5</sup>	n.a. <sup>5</sup>						
Benzo[a]pyrene <sup>5</sup>	n.a. <sup>5</sup>						
Benzo[b,k]fluoranthene <sup>5</sup>	n.a. <sup>5</sup>						
Benzo[e]pyrene <sup>5</sup>	n.a. <sup>5</sup>						
Benzo[g,h,i]perylene	30						
Biphenyl	50						
Chrysene <sup>5</sup>	n.a. <sup>5</sup>						
Coronene <sup>5</sup>	n.a. <sup>5</sup>						
Cyclopenta[c,d]pyrene <sup>5</sup>	n.a. <sup>5</sup>						
Dibenzo[a,h]anthracene <sup>5</sup>	n.a. <sup>5</sup>						
Fluoranthene <sup>5</sup>	n.a. <sup>5</sup>						
Fluorene	40						
Indeno[1,2,3-c,d]pyrene <sup>5</sup>	n.a. <sup>5</sup>					_	
Naphthalene	40						
Phenanthrene	40						
Pyrene <sup>5</sup>	n.a. <sup>5</sup>						
Sum PAHs <sup>13</sup>	0.05 (as	0.0006 (as	0.08	0.65	9.0	84.6	
	BaP)	BaP)					

#### Table 4 – Remarks

- 1. For aluminium, antimony, arsenic, lead and nickel, background values for adults and children are discriminated.
- 2. Background value of aluminium for adults is between 80 and 180 μg/kg/day.
- 3. Background value of antimony for adults is between 0.018 and 0.48 µg/kg/day.
- 4. Background value of arsenic for children is between 0.4 and 0.7  $\mu$ g/kg/day.
- 5. TDI is not available (= n.a.).

- Chromium III must be differentiated into soluble (TDI=5 μg/kg/day) and insoluble (TDI=5000 μg/kg/day).
- 7. Background chromium VI between 5.7×10<sup>-6</sup> 4.3×10<sup>-4</sup> μg/kg/day.
- 8. Non industrial setting.
- 9. Sum of organotins: dibutyltin (DBT), monobutyltin (MBT), monooctyltin (MOT), tributyltin (TBT).
- 10. Alpha-Chlordane = cis-Chlordane.
- 11. Gamma-Chlordane = trans-Chlordane.
- 12. Lindane is technical gamma-HCH.
- 13. Sum of PAHs: acenaphthene, acenaphthylene, anthracene, benzo[a]anthracene, benzo[a]pyrene, benzo[b,k]fluoranthene, benzo[e]pyrene, benzo[g,h,i]perylene, biphenyl, chrysene, coronene, cyclopenta[c,d]pyrene, dibenzo[a,h]anthracene, fluoranthene, fluorine, indeno[1,2,3-c,d]pyrene, naphthalene, phenanthrene, pyrene.
- 14. Provisional value due to limited toxicological data base.
- 15. Sum of aldrin and dieldrin.
- 16. Sum of DDD, DDE and DDT.
- 17. TDI of dimethyl phthalate (DMP) is unknown, as the compound is chemically comparable to diethyl phthalate (DEP), the concentrations of DMP are added to the concentrations of DEP and compared to the TDI of DEP.
- 18. TDI of diisobutyl (DiBP) and dimethylpropyl phthalate (DMPP) are unknown, as these compounds are chemically comparable to di-n-butyl phthalate (DBP), the concentrations of DiBP and DMPP are added to the concentrations of DBP and compared to the TDI of DBP.
- 19. For BDE 47, 100, and 183 no toxicological reference dose (e.g. TDI) is available to compare the exposure with. Therefore, the exposure via intake of house dust is compared to the exposure due to food intake in section 6.12.
- 20. Background exposure to PBDE congeners is based on dietary intake for an adult (A) and a 2-year old child (C) (De Winter-Sorkina *et al.*, 2006).
- 21. No information is available about the intake of BDE209 via food intake, neither is a toxicological reference dose available. Therefore, the exposure via dust of BDE209 can only be compared to the exposure via dust of other PBDE congeners.
- 22. Background value of DEHP for adults is between 3 and 16 μg/kg/day.
- 23. Background value of DEHP for children is between 12 and 26 µg/kg/day.

## 5.2 Levels of substances in particulate matter in air

The amount of inhaled house dust particles is low (< 2%) in comparison to ingested house dust. However, in some cases such as for toxic metals local effects in the lungs may occur. In order to evaluate the possibility of a potential health risk of several metals in air, the highest concentrations in particulate matter in air as reported in selected literature sources were compared to the corresponding TCA-values (chronic limit values for air as mg/m³), see Table 5. Below the TCA no health effects are anticipated even after lifelong exposure. None of the highest concentration in airborne particles exceeds the TCA. For arsenic however the guideline value for air of 6 ng/m³, which will be effective within the EU in

2013, is exceeded, indicating that concentrations of this metal may be undesirably high even though they remain well below the current TCA as used by RIVM.

Table 5. Maximum concentrations of various metals in particulate matter (PM10 or PM2.5) in indoor air.

	Highest conc	Reference air	TCA	Reference
	in air (ng/m <sup>3</sup> )	concentrations	$(ng/m^3)$	TCA
Aluminium	100	(Rasmussen et al., 2007)	unknown	
Arsenic	31	(Oomen et al., 2007)	1000	RIVM 2001
Cadmium	1.3	(Oomen et al., 2007;	5	EU 2000
		Rasmussen et al., 2007)		
Chromium	29	(Oomen et al., 2007)	60000*	RIVM 2001
Cobalt	25	(Oomen et al., 2007)	500	RVIM 2001
Manganese	10	(Rasmussen et al., 2007)	150	WHO 2000
Nickel	20	(Oomen et al., 2007)	50	RIVM 2001
Lead	51	(Oomen et al., 2007)	500	WHO 2000
Zinc	606	(Oomen et al., 2007)	unknown	

<sup>\*</sup> TCA for insoluble Cr-III

## 5.3 Substances without potential human health risk

The majority of the substances present in house dust is without potential human health risk. Compounds without human health risk are: antimony, barium, beryllium, cobalt, magnesium, mercury, molybdenum, selenium, silver, strontium, tellurium, thallium, tin, titanium, uranium, vanadium, organotin compounds, 2,4-dichlorophenoxyacetic acid, alachlor, aldrin, alpha-chlordane, alpha-HCH, atrazine, azinphos methyl, beta-HCH, carbaryl, chloroprofam, chlorpyrifos, DDD, DDE, DDT, diazinon, dicamba, dieldrin, ethyl parathion, gamma-chlordane, glyphosate, heptachlor, lindane, malathion, mecoprop, methamidophos, methyl parathion, metolachlor, pendimethalin, permethrin, phosmet, picloram, resmethrin, tetramethrin, trifluralin, BBP, DEP, DiBP, DIDP, DINP, DMP, DMPP, DBP.

## 5.4 Substances with potential human health risk

Based on the calculated mean and maximum ingestion rates, exposure to the following substances in house dust may lead to potential human health effects. Potential human risk is subdivided into risk for adults and risk for children.

Substances in house dust with potential risk for <u>adults</u>: cadmium, chromium, copper, lead, manganese, BFRs, and sum PAHs.

Substances in house dust with potential risk for <u>children</u>: aluminium, arsenic, cadmium, chromium, copper, lead, manganese, nickel, zinc, DEHP, BFRs, and sum PAHs.

Table 6 shows the substances in house dust with a potential risk for adults and children, based on mean and maximum values. Since the maximum concentration in house dust is not always a realistic value, the P95 is added in the calculations. The highest mean, maximum and P95 of the concentration in house dust are presented. The P95 value was used for the calculation of the risk index and for the contribution of house dust to the TDI as a percentage. Both the P95 risk index and the percentage of TDI were calculated for adults and children separately.

Table 6. Compounds with potential risk for human health

	Concentration in house dust			Risk index		Exposure via	
				based on highest		house dust based	
				P95 conc in house dust		on highest P95 conc <b>compared</b> <b>to TDI</b> 1	
	Highest	Highest	Maximum				
	mean	P95					
	(mg/kg)	(mg/kg)	(mg/kg)	(-)	(-)	(-)	(-)
				Adult	Child	Adult	Child
Compound							
Aluminium	24281	44225	51100	0.28	0.79	0.042	0.39
Arsenic	70	$114^2$	192	$0.38^{2}$	<b>1.46</b> <sup>2</sup>	$0.081^2$	$0.76^2$
Cadmium	13	17.32	220	0.92	1.13	0.025	0.23
Chromium III soluble	159	191.8	5440	0.12	0.35	0.027	0.26
Chromium III	159	191.8	5440	0.00	0.00	0.00	0.00
insoluble							
Copper	261	489	12540	0.73	0.76	0.004	0.039
Lead	1200	1312	37000	0.57	2.99	0.26	2.4
Manganese	260	407	9410	0.81	0.83	0.002	0.017
Nickel	47	116	243	0.41	0.88	0.008	0.078
Zinc <sup>3</sup>	628	1570	30600	0.70	0.72	0.002	0.021
Di(2-ethylhexyl)-	3214	7063	40459	0.42	1.46	0.10	0.94
phthalate							
(DEHP)	0.20	12	5.4	0.20	1.0	0.10	1.7
Sum PAHs <sup>4</sup>	0.29	13	54	0.20	1.8	0.18	1.7
BDE99 – EU <sup>5</sup>	$0.022^5$	$0.15^6$	7.8	0.8	5.0	0.41	3.8

#### Table 6 – Remarks

1. Exposure via house dust compared to TDI calculated as a factor: exposure via house dust (P95) (μg/kg bw/day)

TDI (µg/kg bw/day)

- 2. P90 value, calculation of risk index and % of TDI based on P90 value.
- 3. Non industrial setting
- 4. Surrogate approach (also referred to the 'indicator approach').
- 5. For BDE99 no TDI is available. Instead a maximal allowable intake level derived by De Winter-Sorkina *et al.* (IPCS (International Programme on Chemical Safety), 2001) is employed.
- 6. Geometric mean, and 95<sup>th</sup> percentile of pooled data of (Ibarra *et al.*, 2006; Knoth, 2003; Pless-Mulloli *et al.*, 2006; De Boer, 2007; Santillo *et al.*, 2003)

34

# 6 Discussion substances with potential human health risk

In the present chapter those substances are discussed in more detail for which a potential health risk due to house dust was identified using the highest house dust concentration encountered in literature (chapter 5). In addition, some measures to decrease the exposure to substances via house dust are discussed.

#### 6.1 Aluminium

Table 7. Overview of risk indices for <u>aluminium</u> based on geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is listed in the second columns for an adult and child respectively by comparison to the TDI.

	Ad	lult	Child		
	Risk Index Exposure house		Risk Index	Exposure house	
		dust compared		dust compared	
		to TDI <sup>1</sup>		to TDI <sup>1</sup>	
Mean conc.	0.26	0.023	0.62	0.22	
P95 conc.	0.28	0.042	0.79	0.39	
Max. conc.	0.29	0.050	0.85	0.45	

Table 7 – Remarks

Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.
 Exposure house dust compared to TDI= exposure via house dust (μg/kg bw/day)
 TDI (μg/kg bw/day)

Only one study was found that provides information on the levels of aluminium in house dust. When the P95-concentration or the maximum concentration of aluminium in house dust was used in the calculation of the risk index, the risk index was high but not larger than 1 (0.85 for a child with the maximum concentration in house dust, 0.79 for a child with the 95<sup>th</sup>-percentile data, see Table 7). This indicates only a limited potential health risk. Nevertheless, the contribution of exposure via house dust is not negligible for children (39% of the TDI is filled up by exposure via house dust based on the 95<sup>th</sup>-percentile data). Background exposure to aluminium via food and water intake accounts for about 40% of the TDI for children and 11-24% for adults. Thus, the only study giving information on aluminium exposure through house dust indicates a substantial contribution of this route compared to other sources of exposure. But the relevance of this

one study performed in Ottawa, Canada, for the Dutch situation is unknown. In conclusion, the available information indicates house dust may substantially contribute to total aluminium exposure. It is however not likely that this exposure would lead to human health risks

#### 6.2 Arsenic

Table 8 provides an overview of the risk indices for arsenic. The risk index obtained using the highest concentration in house dust encountered in literature was 2.0 for a child and 0.44 for an adult, whereas the risk indices associated with the highest geometric mean concentration of arsenic in house dust were 1.2 and 0.35, respectively. The background exposure to arsenic for a child already fills up 40-70% of the TDI due to intake by food and water.

Arsenic is a naturally occurring element in the environment that may be released from industrial processes, such as mining activities, metal smelting and burning of fossil fuels. High values of arsenic in house dust are often found near smelters. Yet, also in non-industrial settings the arsenic concentration in house dust is occasionally high. In the Netherlands, high arsenic concentrations may be anticipated in the area of zinc smelters (e.g. Budel-Dorplein). Recent research in 15 houses in Budel-Dorplein determined a arsenic concentration of at maximum 60 mg/kg in house dust, whereas in the control area (Liempde, non-industrial setting) in one house an arsenic concentration in house dust of 74 mg/kg was found (Oomen *et al.*, 2007). This house dust concentration of 74 mg/kg combined with highest background exposure for children (0.7 μg/kg bw/day is used as background exposure of the range 0.4-0.7 μg/kg bw/day) and 0.3 μg/kg bw/day for adults, results in a risk index of 0.35 and 1.2 for an adult and child, respectively. In this calculation the exposure via house dust represents 5 and 49% of the TDI, respectively.

Note that in the earlier research by Oomen *et al.* (2007) a lower dust intake was used in the calculation and exposure was integrated over a lifetime. The conclusion there was that arsenic in house dust from Budel-Dorplein did not cause a potential risk for human health (Oomen *et al.*, 2007).

In conclusion, the present analysis based on literature data indicates substantial exposure to arsenic through house dust. The calculated risk index indicates a potential human health risk for children. For adults background estimated exposure is lower than for children (0.3 versus 0.7  $\mu g/kg/day$ ). Thus the exceedance of the TDI will in most cases be limited to the childhood years. Given its derivation, the TDI for arsenic should be treated as a long-term average, which implies that in principle any temporary exceedance early in life could be compensated by a proportionally lower exposure later in life, as seems to be the case for arsenic in house dust. Nevertheless, given the general picture for arsenic health risks of only a limited margin between actual intakes and levels known to produce toxic



effects when humans are exposed to them chronically, arsenic exposure in general should preferably be as low as possible.

Table 8. Overview of risk indices for <u>arsenic</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the TDI.

oxpodulo lo doc	expectate to december in the ecocita column for an additional of the by			
	A	dult	Child	
	Risk Index	Exposure house	Risk Index	Exposure house
		dust compared		dust compared
		to TDI <sup>1</sup>		to TDI <sup>1</sup>
Mean conc.	0.35	0.05	1.2	0.47
P95 conc.	0.38	0.08	1.5	0.76
Max. conc.	0.44	0.14	2.0	1.3
Highest conc.	0.35	0.05	1.2	0.49
Netherlands <sup>2</sup>				

#### Table 8 – Remarks

- Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.
   Exposure house dust compared to TDI= exposure via house dust (μg/kg bw/day)
- 2. Highest concentration encountered in a study in the Dutch Kempen area (Oomen *et al.*, 2007).

#### 6.3 Cadmium

Important for cadmium is that background exposure due to food and water intake already accounts for 90% of the TDI. This limits the allowable additional exposure via other routes. Thus relatively low levels of cadmium in house dust may already lead to risk indices higher than 1.0, indicating a potential health risk. Occasionally cadmium levels in house dust are so high that they may pose a potential health risk even without taking into account background exposure, see Table 9. In most cases, however, the contribution of house dust to total cadmium exposure is limited. Thus for cadmium, exposure through house dust may give rise to a potential health risk when taking the background into account, but its contribution to total exposure mostly is limited only.

The highest cadmium concentration recently encountered in house dust in the Netherlands was 22.8 mg/kg in Budel-Dorplein, an area known to be historically contaminated with cadmium (Oomen *et al.*, 2007). The cadmium concentration in all other houses (n=45) was considerably lower (< 7 mg/kg). This highest concentration in the Netherlands is, according to the present method, associated with a risk index of 0.93 and 1.2 for an adult

and child, respectively (accounting for 3.2% and 30 % of the TDI, respectively), see Table 9. This confirms the picture obtained form literature.

Cadmium is an extremely potent nephrotoxicant. This toxicity develops over a time period of decades of continuing low exposure, with the kidney cortex as the target tissue. Thus a risk index for a child greater than 1 does not necessarily mean an actual health risk during this period of life. Exposure to cadmium should be evaluated integrally over the childhood and adult years. When modelling exposure to the highest cadmium concentration in house dust in Budel-Dorplein (22.8 mg/kg) during 7 childhood years, 11 transition years and 62 adult years, the critical cadmium levels in the kidney cortex were calculated not to be reached, and therefore no health risk was anticipated (Oomen *et al.*, 2007). Despite this absence of an actual health risk, elevated exposure to cadmium is undesirable in principle and a policy of reducing its exposure via food and other routes is followed by health authorities. Thus levels in house dust should also be as low as possible.

Table 9. Overview of risk indices for <u>cadmium</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in

the second column for an adult and child by comparison to the TDI.

	Ad	Adult		ild
	Risk Index	Exposure house	Risk Index	Exposure house
		dust compared		dust compared
		to TDI <sup>1</sup>		to TDI <sup>1</sup>
Mean conc.	0.92	0.02	1.1	0.17
P95 conc.	0.92	0.03	1.1	0.23
Max. conc.	1.2	0.31	3.8	2.9
Highest conc. Netherlands <sup>2</sup>	0.93	0.03	1.2	0.30
Netherlands <sup>2</sup>				

#### Table 9 – Remarks

- Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.
   Exposure house dust compared to TDI= exposure via house dust (μg/kg bw/day)

  TDI (μg/kg bw/day)
- 2. Highest concentration encountered in a study in the Dutch Kempen area (Oomen *et al.*, 2007).

A further issue with cadmium is its inhalatory carcinogenicity as evident from in studies in industrial workers with high exposures. For cadmium a toxicological reference value for air of 5 ng/m³ has been derived as protective against both its nephrotoxicity and its carcinogenicity (Oomen *et al.*, 2007). The highest cadmium concentration in air samples in Budel-Dorplein, Maarheeze and Liempde in the Netherlands was 1.4 ng/m³ (n=45) (Oomen *et al.*, 2007). It can be calculated that theoretical exposure to this highest cadmium concentration in air during an entire life, would lead to an extra cancer risk of  $3.2-5.8 \times 10^{-6}$  (32 to 58 additional cases of cancer per 10 million lifelong exposed). Based

on the available evidence it is concluded that cadmium exposure via dust particles in the air does not exceed the toxicological reference value of 5 ng/m<sup>3</sup>.

#### 6.4 Chromium

From a toxicological point of view trivalent and hexavalent chromium should be differentiated. Hexavalent chromium is much more toxic than trivalent chromium, having a much higher toxic potential for different toxicological endpoints and most importantly, unlike the trivalent form, presenting a genotoxic and carcinogenic risk. Soluble trivalent chromium in turn is more toxic than insoluble trivalent chromium, which is reflected by their respective TDIs of 5 and 5000 µg/kg/day.

Available data on chromium concentrations in house dust represent total chromium, i.e. without specification of valence and solubility. No general information is available on the contribution of hexavalent chromium in house dust. A priori it is plausible that chromium in house dust is trivalent chromium due to the chemical instability of hexavalent chromium. However, as indicated, this assumption cannot be founded by actual data. Trivalent chromium is expected to be present predominantly as insoluble compound (in parallel to soil in which the insoluble forms as carbonate and oxide dominate, chromium dissolves only when complexation is possible).

Assuming chromium to be present in house dust predominantly as insoluble trivalent ion, all risk indices are close to zero, indicating absence of a potential risk.

### 6.5 Copper

Copper is an essential trace nutrient to humans. Nevertheless high concentrations of copper can be toxic. Based on the maximum value in house dust, copper can cause a potential risk for children (risk index 0.83 and 1.7 for an adult and child, respectively, see Table 10). When calculations were performed using the P95 or the highest geometric mean, no potential risk was found. The highest copper concentration in house dust encountered in literature (12540 mg/kg) was much higher than the 95<sup>th</sup>-percentile concentration (489 mg/kg) and the geometric mean (261 mg/kg). This indicates that the highest value may be an extreme situation. It is therefore unlikely that background exposure and exposure via house dust to copper would actually result in a potential risk for human health.

Crucially, in general copper exposure via house dust accounts for only a small percentage of the TDI (about 0.4% for a child at the P95-exposure level). About 72% of the TDI is filled up by background exposure via food and water intake. Hence, house dust is in most cases only a minor exposure route for copper.

Table 10. Overview of risk indices for <u>copper</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the TDI.

	Adult		Child	
	Risk Index	Exposure house dust compared to TDI <sup>1</sup>	Risk Index	Exposure house dust compared to TDI <sup>1</sup>
Mean conc.	0.73	0.002	0.74	0.021
P95 conc.	0.73	0.004	0.76	0.039
Max. conc.	0.83	0.11	1.7	1.0

#### Table 10 – Remarks

1. Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.

Exposure house dust compared to TDI=  $\frac{\text{exposure via house dust (<math>\mu g/kg \text{ bw/day})}}{\text{TDI (}\mu g/kg \text{ bw/day)}}$ 

#### 6.6 Lead

As can be seen in Table 11, the TDI of lead can be exceeded considerably when both background exposure and exposure to house dust are taken into account. House dust exposure on its own already exceeds the TDI on a regular basis, as becomes apparent from the risk index of 2.7 for children based on the highest geometric mean value of lead in house dust found in literature.

As is known from the extensive literature on lead health effects, children are a vulnerable group for lead toxicity and the TDI for lead has been derived from data on health effects in this particular group. The basic principle in this derivation is that any increase in blood lead levels in children is unwanted from a toxicological point of view given the high neurotoxic potential of lead as demonstrated in numerous human studies. Thus this TDI does not incorporate the usual margins introduced by uncertainty factors but is based on actual data from the sensitive subgroup in the population. This implies that exposure to lead cannot be integrated over childhood and adult years, and thus that exceedance of the TDI for children represents a potential health risk. As can be seen from Table 12, exposure via house dust results in potential health risks on a regular basis, and house dust is an important exposure route in many cases. It is therefore concluded that lead in house dust results in potential health risks on a regular basis.

In a recent study by RIVM (Oomen *et al.*, 2007) lead concentrations in house dust from 45 houses were determined. Both in the near vicinity of a zinc smelter and in the control area lead concentrations in house dust were occasionally high (up to 2560 mg/kg). The latter lead concentration in house dust is associated with a risk index of 0.81 for an adult and 5.2 for a child, representing 51% and 474% of the TDI of lead without background.

The high lead concentrations in house dust were assumed to be mainly related to certain hobbies or behaviours (soldering, hunting), and in most cases no clear relationship with the environment could be found (Oomen *et al.*, 2007). However, other studies have shown that also outdoor soil may cause high lead levels in indoor house dust, due to the large fraction of soil components in house dust (Gulson *et al.*, 1995).

Table 11. Overview of risk indices for <u>lead</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the TDI.

	Adult		Child	
	Risk Index	Exposure house	Risk Index	Exposure house
		dust compared		dust compared
		to TDI <sup>1</sup>		to TDI <sup>1</sup>
Mean conc.	0.54	0.24	2.7	2.2
P95 conc.	0.57	0.26	2.9	2.4
Max. conc.	7.7	7.3	69	69
Highest conc.	0.81	0.51	5.2	4.7
Netherlands <sup>2</sup>				

#### Table 11 – Remarks

- Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.
   Exposure house dust compared to TDI= exposure via house dust (μg/kg bw/day)
- 2. Highest concentration encountered in a study in the Dutch Kempen area (Oomen *et al.*, 2007).

For lead it should be noted that for the calculation of the above risk indices a background exposure was used reflecting the intake via food and water consumption and via soil ingestion, i.e. exclusive of house dust. This estimate was based on a document from the Dutch Health Council (Dutch Health Council (Gezondheidsraad), 1997) in which estimates were developed for the daily intake of lead by the Dutch population via different routes. For a child a maximum intake of 1.33 µg/kg bw/day was estimated for food and water intake. For an adult this was estimated at 0.86 µg/kg bw/day. Soil and dust intake together was estimated to account for 1.0 µg/kg bw/day for a child. A child would have about half this amount via soil and dust. Assuming that lead exposures via soil and dust intake contribute approximately equally (0.5 µg/kg bw/day) a background exposure of 1.83 µg/kg bw/day is derived for a child and 1.11 µg/kg bw/day for an adult.

### 6.7 Manganese

Manganese is an essential trace nutrient to humans. Manganese is present in many foods, including grains and cereals, and is found in high concentrations in many foods, such as tea. Based on the maximum value in house dust, manganese was identified as a potential risk for children (risk index 1.2, see Table 12). When calculations were performed using the P95 or the highest geometric mean, no potential risk was found (risk index 0.83). As can be seen in Table 6, there is a large difference between the maximum concentration (9410 mg/kg) and the P95-concentration (407 mg/kg). The maximum value is assumed to occur in extreme conditions. Only two references with data on manganese levels in house dust were found. Although the two references described similar concentration in house dust, the variation in manganese levels in house dust and the relevance of these data for the Dutch situation are unknown.

Importantly, the contribution of house dust accounts for only a few percent of the TDI in most cases (1.7% for the P95-exposure values for children). It is therefore concluded that it is unlikely that manganese in house dust in will increase background exposure to the extent of posing an actual risk for human health.

Table 12. Overview of risk indices for <u>manganese</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the TDI.

	Adult		Child	
	Risk Index	Exposure house dust compared to TDI <sup>1</sup>	Risk Index	Exposure house dust compared to TDI <sup>1</sup>
Mean conc.	0.81	0.001	0.82	0.011
P95 conc.	0.81	0.002	0.83	0.017
Max. conc.	0.85	0.042	1.2	0.39

Table 12 – Remarks

1. Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.

Exposure house dust compared to TDI=  $\frac{\text{exposure via house dust } (\mu g/kg \text{ bw/day})}{\text{TDI } (\mu g/kg \text{ bw/day})}$ 

#### 6.8 Nickel

The risk index for nickel does not exceed 1, not even for the highest encountered house dust concentration in literature. However, the risk index was very close to 1 (0.96 for

highest dust concentration for children, see Table 13). Background exposure of a child already accounts for 80% of the TDI. The contribution of house dust to the total exposure is usually low (7.8% of the TDI is filled up for the P95-exposure data). The risk index for adults is even for the maximum concentration in house dust less than 0.5 (0.42), indicating that a health risk is not anticipated.

Table 13. Overview of risk indices for <u>nickel</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the TDI.

expectate to decertable in the ecocita column for an au			it ana onia ay ooni		
	Ac	Adult		Child	
	Risk Index	Exposure house	Risk Index	Exposure house	
		dust compared		dust compared	
		to TDI <sup>1</sup>		to TDI <sup>1</sup>	
Mean conc.	0.40	0.003	0.83	0.031	
P95 conc.	0.41	0.008	0.88	0.078	
Max. conc.	0.42	0.017	0.96	0.16	

Table 13 – Remarks

Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.
 Exposure house dust compared to TDI= exposure via house dust (μg/kg bw/day)
 TDI (μg/kg bw/day)

#### **6.9 Zinc**

Zinc is one of the most common elements in the earth's crust. Zinc is found in air, soil, and water and is present in all foods. Zinc enters air, water, and soil as a result of both natural processes and human activities. The latter involve mining, purifying of zinc, lead, and cadmium ores, steel production, coal burning, and burning of wastes.

Exposure to zinc via house dust usually accounts for a minor part of the total daily exposure only (2% of the TDI for children is filled up for the P95-concentration in house dust, see Table 14). The background exposure via food already accounts for 70% of the TDI. The risk index is only greater than 1 for children at the highest encountered house dust concentration of 30600 mg/kg. This appears to be an exceptionally high concentration, as the concentration at the 95<sup>th</sup>-percentile is 1570 mg/kg. In a recent study in the Netherlands in 45 houses, including 15 houses in the direct neighbourhood of a zinc smelter, the highest zinc concentration in house dust was 1724 mg/kg. Hence, it is concluded that it is highly unlikely that zinc in house dust poses a potential health risk to humans.

Table 14. Overview of risk indices for <u>zinc</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the TDI.

	Adult		Child	
	Risk Index	Exposure house	Risk Index	Exposure house
		dust compared		dust compared
		to TDI <sup>1</sup>		to TDI <sup>1</sup>
Mean conc.	0.70	0.001	0.71	0.008
P95 conc.	0.70	0.002	0.72	0.021
Max. conc.	0.74	0.044	1.1	0.41
Highest conc. Netherlands <sup>2</sup>	0.70	0.003	0.72	0.023
Netherlands <sup>2</sup>				

#### Table 14 – Remarks

- Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.
   Exposure house dust compared to TDI= exposure via house dust (μg/kg bw/day)
- 2. Highest concentration encountered in a study in the Dutch Kempen area (Oomen *et al.*, 2007).

### 6.10 Di(2-ethylhexyl)phthalate (DEHP)

DEHP is used as a plasticiser in PVC. Because of its toxic potential specifically for human reproductive development (hormonal disruption) DEHP along with several other phthalates has been banned from use in toys and childcare articles within the EU (Directive 2005/84/EC). For other applications these compounds remain in use, including in food contact materials and various consumer products such as carpets and upholstery.

Background exposure to DEHP via food and water intake is estimated at 12-26 μg/kg bw/day for a child and 3-16 μg/kg bw/day for an adult (EFSA, 2005). The background exposure already fills up the TDI of 50.0 μg/kg bw/day to 24-52% for a child, and 6-32% for an adult. In addition to background exposure, house dust is an important exposure route to DEHP. The risk index is exceeded 5.9-fold for the highest house dust concentration described in literature for children, with house dust responsible for a 5.4-fold exceedance of the TDI, see Table 15. Already the highest geometric mean concentration of DEHP in house dust results in a risk index near 1 for children (0.95; of which 0.43 is due to exposure to house dust only). The risk index for adults is not exceeded. Given the toxicological basis for the TDI for DEHP with disruption of reproductive development in males as the critical effect, lower levels of exposure later in life do not compensate for exceedance of the TDI during child age. Hence, exposure to

44

DEHP should not be integrated over life, and risk indices for children should be evaluated as such. In conclusion, DEHP poses a human health risk, both due to the background exposure and due to house dust intake.

It is unlikely that the other phthalates result in potential health risks, although the risk index for DBP in combination with DMPP and DiBP is rather high for the highest concentration found in literature (0.73 for children). Also the risk index for the highest concentration found in literature of BBP is not negligible (0.63 for children).

Table 15. Overview of risk indices for <u>DEHP</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the TDI.

•	Adult		Child	
	Risk Index	Exposure house	Risk Index	Exposure house
		dust compared		dust compared
		to TDI <sup>1</sup>		to TDI <sup>1</sup>
Mean conc.	0.37	0.05	0.95	0.43
P95 conc.	0.42	0.10	1.5	0.94
Max. conc.	0.90	0.58	5.9	5.4

Table 15 – Remarks

Exposure via house dust compared to TDI calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the TDI.
 Exposure house dust compared to TDI= exposure via house dust (μg/kg bw/day)
 TDI (μg/kg bw/day)

#### 6.11 Sum PAHs

A number of PAHs, including benzo(a)pyrene (BaP), are genotoxic carcinogens. In the present report the so-called *indicator approach* (also denominated as the 'surrogate approach') was used for PAHs, which involves using benzo(a)pyrene as the marker for the total mixture, both in quantifying risk and exposure. Although the use of the more complex 'potency approach' (in which for individual PAHs potency factors applied to calculate BaP-equivalents) is in principle more appropriate for environmental contaminations (see RIVM, 2001), within the inherent uncertainty boundaries of the present calculation of Risk Indices, we deemed the use of the more simple indicator approach acceptable.

For the total mixture of PAHs the Maximum Permissible Risk (MPR), which according to Dutch environmental policy equals the risk-specific dose for one in ten thousand per lifetime, is 50 ng BaP/kg bw/day (this figure applies to BaP as the marker for the PAH mixture as a whole). Background exposure to PAHs is low compared to this MPR

(6 ng BaP/kg bw/day for an adult and 15 ng BaP/kg bw/day for a child). Remarkably the MPR is sometimes exceeded due to exposure via house dust. For the 95<sup>th</sup> percentile data, the risk index is 0.31 and 2.0 for adults and children, respectively, see Table 16. The risk index for the geometric mean concentration of the sum of PAHs in house dust is 0.12 and 0.34 for adults and children, respectively. In conclusion, in most cases the PAH contents of house dust does not result in a health risk higher than the MPR. However for high PAH concentrations (e.g. 95<sup>th</sup> percentile) exposure via house dust is greater than via food, leading even to exceedance of the MPR. Given the general policy of reducing exposures to the genotoxic and carcinogenic PAH via food to levels as low as reasonably possible, this high exposure via house dust is significant.

Indoor sources of PAHs include cooking, heating, cigarette-smoking, wood burning, candle burning, and incense burning. Outdoor sources include vehicle exhaust and industrial processes such as aluminium smelting, coke production, and petroleum refining (Maertens *et al.*, 2004). Maertens *et al.* reviewed the data of 18 studies and concluded that an urban location and the presence of cigarette smokers increased the PAH content of house dust (Maertens *et al.*, 2004). Yet, correlations were weak, indicating that other factors (e.g. flooring type, season, deposition rate, ventilation, social-economic status) may affect the PAH content as well.

Table 16. Overview of risk indices for <u>sum of PAHs</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is accounted for). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the MPR<sup>2</sup>.

	Adult		Child	
	Risk Index	Exposure house dust compared to MPR <sup>2</sup>	Risk Index	Exposure house dust compared to MPR <sup>2</sup>
Mean conc.	0.12	0.004	0.34	0.04
P95 conc.	0.31	0.18	2.0	1.7
Max. conc.	0.89	0.77	7.5	7.2

#### Table 16 – Remarks

- 1. Calculation for the sum of PAHs is based on a so-called 'surrogate approach'. The benzo[a]pyrene concentration is considered to be representative for the entire PAH concentration, but is compared to a 10-times lower toxicological reference value to compensate for the other PAHs.
- 2. For genotoxic carcinogens no TDIs are derived but the denomination 'MPR' is used, defined within Dutch environmental policy as an extra cancer risk of 1 in 10.000 people based on lifetime exposure. Exposure via house dust compared to MTR calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the MPR.

Exposure house dust compared to MPR=  $\frac{\text{exposure via house dust } (\mu g/kg \text{ bw/day})}{\text{MPR } (\mu g/kg \text{ bw/day})}$ 

#### 6.12 Brominated flame retardants

#### **6.12.1** Toxicological reference values

A major issue with brominated flame retardants is that internationally accepted TDI values for either individual congeners or the sum of brominated flame retardants are not yet available (De Winter-Sorkina *et al.*, 2006). To compensate for this, De Winter-Sorkina *et al.* derived a (provisional) maximal allowable intake level for one congener, BDE99, of 0.26 ng/kg bw/day (De Winter-Sorkina *et al.*, 2006; Bakker *et al.*, in press), which is also used here. In the same report also the median of the averaged life-long daily dietary intake of BDE99 is estimated at 0.11 ng/kg bw/day for the Dutch population (0.23 and 0.10 ng/kg bw/day for a 2-year old child and an adult, respectively). As indication of the distribution of the exposure to BDE99 via food, the 99<sup>th</sup>-percentile of the averaged lifelong daily dietary intake to BDE99 is given, amounting to 0.24 ng/kg bw/day.

#### 6.12.2 Ban on several brominated flame retardants

Not all PBDEs are still in use, although differences between countries exist. The penta-BDE technical product (including BDE 99) was voluntarily withdrawn from use in the European Union during the last 10 years. Use of penta-BDE and octa-BDE technical products in all applications for the European Union market has officially been banned since August 2004. The use of penta-BDE, octa-BDE and PBBs in new electrical and electronic equipment has been banned since July 2006 (De Winter-Sorkina *et al.*, 2006). However, hexa-BDE (e.g. BDE 153 and 154) and deca-BDE (e.g. BDE 209) and other groups of brominated flame retardants are still in use, with especially a high production of deca-BDE (e.g. BDE 209) (De Boer and Wells, 2006). It has been postulated that due to the European ban on several PBDEs, concentrations of most PBDEs in house dust in this part of the world might be significantly lower than concentrations in the United States (Harrad *et al.*, in press). However, this is still under discussion.

#### 6.12.3 Possibility of a potential health risk for BDE99

A potential health risk due to exposure via house dust is identified for the only congener for which a provisional toxicological reference value is available (BDE99). This is shown by the calculated risk indices, which are for adults and children 0.45 and 1.5 respectively for the geometric mean concentration data in house dust, 0.8 and 5.0 for 95<sup>th</sup>-percentile data, and 22 and 204 for the maximum concentration data encountered in the EU (see Table 17). As background exposure the estimated median daily dietary intake levels for a child (0.23 ng/kg bw/day) and an adult (0.10 ng/kg bw/day) were used. The contribution of house dust to the exposure to BDE99 is considerable, especially for children. For example, 6% and 57% of the allowable daily intake level is filled up by dust intake for adults and children based on the EU geometric mean data.

For comparison to the maximum allowable intake level, exposure over an entire lifetime should be considered. As Table 17 shows, the expected exposure due to dust (including

food) intake is expected to exceed the maximum allowable intake level on a regularly basis both for adults and children. In addition, a pilot study on the release of PBDEs in a system that simulates the physicochemical conditions of the human gastrointestinal tract suggests that PBDEs are easily released from the dust into the gastrointestinal fluids and can become available for intestinal uptake (De Boer, 2007). This suggests that the bioavailability of PBDEs in house dust may be high. This leads to the conclusion of a potential health risk due to BDE99.

Table 17. Overview of risk indices for <u>BDE99 in Europe</u> based on the highest encountered geometric mean, P95 and maximum concentration data encountered in house dust (background exposure is based on estimated median daily dietary intake levels for children and adults). The contribution of the exposure via house dust to the total exposure is described in the second column for an adult and child by comparison to the (provisional) maximal allowable intake level.

	Ad	Adult		nild
	Risk Index	Exposure house	Risk Index	Exposure house
		dust compared		dust compared
		to maximal		to maximal
		allowable intake		allowable intake
		level <sup>1</sup>		level <sup>1</sup>
Mean conc.	0.45	0.06	1.5	0.57
P95 conc.	0.8	0.41	5.0	3.8
Max. conc.	22	22	204	203

#### Table 17 – Remarks

1. A (provisional) maximal allowable intake level is used because no TDI-value for PBDEs has been derived up till now. Exposure via house dust compared to this (provisional) maximal allowable intake level is calculated as a factor. When the factor is greater than 1, the exposure route via house dust on its own exceeds the intake level.

Exposure house dust compared to intake level=  $\frac{\text{exposure via house dust (}\mu\text{g/kg bw/day)}}{\text{maximal allowable intake level (}\mu\text{g/kg bw/day)}}$ 

#### 6.12.4 Comparison intake via house dust and food

For several other PBDE-congeners than BDE99 for which reliable information on levels in house dust is available, no toxicological reference values are available. Therefore, we only compare exposure to these congeners via intake of house dust to exposure via food. The contribution of house dust seems to be greater for BDE99 than for BDE47, BDE100 and BDE183, see Table 18. Only for high concentrations in dust, intake of BDE47 and BDE100 via this route dominates over the median dietary intake, at least within the EU.

48



Table 18. Ratio of various PBDE-congeners between estimated intake via house dust and food. Ratios are calculated based on geometric mean and 95<sup>th</sup>-percentile ingestion levels via house dust and median dietary intake levels for a child and an adult. When the ratio is greater than 1, intake via house dust dominates over the median intake via food.

	Ratio intake via house dust (unitless)			
	7.5	intake via		7
	Mean ingestion	Mean ingestion	P95 ingestion	P95 ingestion
	adult	child	adult	child
BDE 47 - EU	0.06	0.13	1.2	2.3
BDE 99 - EU	0.16	0.64	1.1	4.3
BDE 99 – US <sup>1</sup>	16	67		
BDE 100 - EU	0.10	0.35	1.7	5.8
BDE 183 - EU	0.02	0.07	0.13	0.46

#### Table 18 – Remarks

1. Levels of PBDE in house dust are much higher (depending on the congener, in most cases at least tenfold) in the United States compared to the European Union.

#### 6.12.5 Comparison of exposure BDE209 via house dust to other congeners

For BDE209 neither a toxicological reference value nor reliable information exposure via food consumption are available. The latter is due to analytical problems related with the analysis of BDE209 in food samples, mainly caused by high background levels (De Boer *et al.*, 2006). The concentrations of BDE209 in house dust are much higher than in food so that these data are assumed to be reliable. Given this, it seems worthwhile to compare exposure of BDE209 via house dust to exposure to other PBDEs via this route. Based on the geometric mean concentrations of the European Union data, the concentration of BDE209 in house dust is higher by factors of 31, 38, 88, and 94 respectively than the concentrations of BDE47, BDE99, BDE100, and BDE183.

This indicates that human exposure of BDE209 via house dust may be (much) greater than via food intake. In addition, when looking at total PBDE exposure via house dust, the major fraction is BDE209. In order to arrive at more definitive conclusions, a toxicological reference value for BDE209 in particular is needed. However, for a full evaluation of the risks of BDEs in general, also toxicity reference values for the other congeners are necessary too. To derive toxicological reference values for BDEs, additional toxicity studies are urgently required.

#### 6.12.6 Other brominated flame retardants

A further brominated flame retardant, tetrabromobisphenol-A (TBBP-A), was not found toxic for rodents in dose levels up to 1000-10000 mg/kg/day (EU RAR, 2003)(De Winter-Sorkina *et al.*, 2006), so this substance does not seem to be of concern for human health.

#### 6.12.7 Non-brominated flame retardants

It should be noted that other flame retardants are emerging, some of which are similarly persistent and bioaccumulative as PBDEs. Compounds such as Dechlorane-plus (a chlorinated flame retardant) (Zhu *et al.*, 2007) and fluor-containing flame retardant are gaining attention as substances possibly posing a human health risk. However, at the moment too little information about these substances is available to be able to draw any conclusions. It is recommended to pay attention to these substances and assess their potential health risk in the near future.

# 6.13 Measures that can decrease the exposure to substances in house dust

A simple measure to decrease the exposure to substances in house dust is cleaning. But only few scientific papers address this issue, making it difficult to draw general conclusions. Some research has focussed on the effect of cleaning on the exposure to lead in house dust. Yinn et al. showed a significant reduction in lead loading on soiled carpets after vacuum cleaning (New Yersey, US) (Yiin et al., 2002). They also performed a randomized study to determine whether a conventional vacuum cleaner could achieve cleaning-results comparable with those of a high-efficiency particulate air filtered vacuum cleaner. They found no differences in lead loading reduction between the two vacuum cleaners, from which they concluded that both vacuum cleaners showed a lead loading reduction. In contrast to this finding, Paustenbach et al. showed that most conventional vacuum cleaners do not trap small particles(< 20 μm) but simply re-emit them into the air (references in (Paustenbach et al., 1997)). Therefore, when considering cleaning as a measure to decrease exposure to substances, Paustenbach et al. advised to do wet cleaning (e.g. mopping, washing). In addition, the Australian EPA advises to clean wet with high phosphate detergents as a measure to decrease exposure to lead via house dust (http://www.epa.nsw.gov.au/leadsafe). It is also recommended to do the wet mopping at least an hour after vacuuming so that dust particles can settle. Based on these studies, wet cleaning is preferred over vacuuming to decrease the exposure to substances present in house dust.

Campbell *et al.* measured dust and blood lead levels in children to monitor the effectiveness of professional home cleaning (Philadelphia, US) (Campbell *et al.*, 2003). They found lower blood lead levels among children in homes with low precleaning dust lead levels, compared with the blood lead levels of children who lived in high-exposure homes, indicating that house dust is an important exposure route for lead. In addition, professional cleaning produced immediate reductions of 36% in dust lead levels. However, lead dust levels rebounded to levels before professional cleaning within three to six months. Thus, regular cleaning will be required if blood lead or dust lead levels are to be reduced and sustained at low levels.

This re-increase in substance levels in dust will occur primarily if the source of the substances in house dust remains present. Where sources can be identified and removed exposure via house dust is structurally decreased. Examples of such sources are mainly known for lead: soldering, presence of lead bullets in the home for hunting purposes, leaded paint (a major problem in older houses in the US). Furthermore, habits such as smoking may increase the amount of several metals and PAHs in the home substantially.

In conclusion, if possible the source of the substances in house dust should be identified and measures should be taken to prevent further contamination in the future. Dust control may be effective if performed properly (preferentially by wet cleaning) and regularly. Under controllend circumstances measures could be evaluated after some time by analysis of dust samples as resurgence to former substance concentrations may occur.

### 7 Conclusions

The present report assesses the substances for which a potential human health risk due to exposure to house dust is possible. The research was limited to metals, organotin compounds, phthalates, brominated flame retardants, pesticides, and polycyclic aromatic hydrocarbons. We estimated exposure levels based on house dust data obtained from literature, also taking into account exposure via other routes (background). Subsequently we compared exposure to toxicological reference values by calculating risk indices (exposure divided by tolerable daily intakes). For most substances a human health risk is not anticipated, even with the conservative assumptions made.

The compounds for which a health risk due to exposure to house dust may be possible are: lead, di(2-ethylhexyl)-phthalate (DEHP), followed by arsenic, cadmium, PAHs and polybrominated diphenyl ethers (PBDEs). Of the PBDEs a health risk could only be identified for congener 99, as for this congener a provisionally maximum allowable intake level was available. Further research is recommended to evaluate the toxicity of the other PBDEs in general. In the present framework, a toxicological reference value is especially needed for BDE209 as the major fraction of the exposure to PBDEs via house dust consists of BDE209, whereas the contribution of food to the total exposure is expected to be small. The other BDE congeners may also result in potential health risks but this is unknown due to the lack of toxicity data, but BDE99 indicates that the exposure to this congener is borderline to its human health exposure limit value (Bakker *et al.*, in press). However, for several other congeners the contribution of food to the total exposure is expected to be larger than the contribution of house dust. Yet, it is important to assess risks to substances through various exposure routes so that health risks can be identified and measures can be taken effectively.

In Table 19 we give qualitative indications of these potential risks based on further analysis. The most important issues with these substances are briefly discussed below.

In conclusion, the substances for which a potential health risk due to exposure via house dust is expected to occur most frequently are <u>lead</u> and <u>DEHP</u>, followed by <u>arsenic</u>, <u>cadmium</u>, <u>sum PAHs</u>, and <u>PBDEs</u> (health risk identified for BDE99; risk assessment was not possible for other BDE-congeners due to lack of toxicity data).

Table 19. Qualitative indications on the substances for which there may be a health risk due to exposure via house dust.

Compound	Potential risk for human health <sup>1</sup>
Aluminium	Unlikely
Arsenic	Likely (but at high end of exposure only)
Cadmium	Likely (but at high end of exposure only)
Chromium	Unlikely
Copper	Highly unlikely
Lead	Likely
Manganese	Highly unlikely
Nickel	Highly unlikely
Zinc	Highly unlikely
Di(2-ethylhexyl)-phthalate (DEHP)	Likely
Sum PAHs	Likely (but at high end of exposure only)
BDE99	Likely (but at high end of exposure only)

#### Table 19 – Remarks

1. A potential health risk is considered to be present if the background exposure and the exposure via house dust are greater than the toxicologically derived reference level expressed as the TDI.

**Aluminium.** Calculated risk indices remained below 1.0, even for children at the maximum aluminium concentration in house dust. Aluminium concentrations in house dust were reported in one study only; no information is available for the Dutch situation. Although contribution of house dust to total aluminium exposure can be considerable (for example 39% of the TDI for P95-exposure data for children), the possibility of a health risk is unlikely. Conclusion, however, is weak due to limited information.

Arsenic. The risk index for a child is often greater than 1, indicating exposure via house dust and background in excess of the TDI. The contribution of house dust to the total exposure is substantial (the highest geometric mean concentration in house dust corresponds with 47% of the TDI). When comparing with the TDI, arsenic exposure should be treated as a long-term average. Due to the lower exposure during adult years the risk index integrated over childhood and adult years is mostly not exceeded. Nevertheless, given the general picture for arsenic health risks of only a limited margin between actual intakes and levels known to produce toxic effects when humans are exposed to them chronically, arsenic exposure in general should preferably be as low as possible.

**Cadmium.** Important for cadmium is that the background exposure to cadmium via food and water intake already accounts for 90% of the TDI. Yet, house dust can contribute considerably in some cases. For example, exposure via house dust at the geometric mean concentration accounts for 17% of the TDI for a child. Given the toxic action of cadmium (induction of nephrotoxicity over a period of many years) exposure during childhood and

adult years should be integrated to assess the possibility of a human health risk. This integration would in many cases result in risk indices < 1. Despite this absence of an actual health risk, elevated exposure to cadmium is undesirable and a policy of reducing its exposure via food and other routes is followed by health authorities. Thus levels in house dust should also be as low as possible.

Chromium. From a toxicological point of view trivalent and hexavalent chromium should be differentiated. Hexavalent chromium is much more toxic than trivalent chromium. Soluble trivalent chromium in turn is more toxic than insoluble trivalent chromium. Available data on chromium concentrations in house dust represent total chromium, i.e. without specification of valence and solubility. A priori it is plausible that chromium in house dust is trivalent chromium due to the chemical instability of hexavalent chromium. Trivalent chromium is expected to be present predominantly as insoluble compound (in parallel to soil in which the insoluble forms as carbonate and oxide dominate, chromium dissolves only when complexation is possible). The estimated risk based on insoluble trivalent chromium in house dust is negligible.

**Copper.** The background exposure to copper via food and water intake already accounts for 72% of the TDI. Exposure via house dust usually represents only a minor exposure route.

Only a potential health risk for children could be calculated with the highest copper concentration found in literature. This highest copper concentration is assumed to be an extreme situation as 95<sup>th</sup>-percentile and geometric mean concentrations were a factor 26 and 48 lower, respectively. Hence, it is highly unlikely that copper in house dust results in a potential health risk.

**Lead.** Exceedance of the TDI for children is unwanted since this population group is especially vulnerable to toxicity caused by lead. Available data indicate that house dust regularly contains lead concentrations that cause a potential health risk for children. For example, at the highest geometric mean concentration the risk index is 2.7, mainly due to exposure to lead via house dust. It is assumed that in many case hobbies or specific behaviour are the cause of high lead concentrations in dust but also lead from outdoor soil may contribute substantially to the high lead levels in dust. In conclusion, a potential health risk due to lead in house dust occurs on a regular basis.

**Manganese.** Only in extreme situations the risk index is exceeded for children (based on the highest manganese concentration described in literature, which was very high in comparison to the P95 value). Background exposure to manganese via food and water intake already accounts for 81% of the TDI. Exposure via house dust usually represents only a minor exposure route. This indicates that it is highly unlikely that manganese in house dust results in a potential health risk.

**Nickel.** Only in extreme situations the risk index can be exceeded (based on the highest nickel concentration described in literature a risk index close to 1 was obtained).

The background exposure to nickel via food and water intake already accounts for 80% of the TDI. Exposure via house dust usually represents only a minor exposure route. This indicates that it is highly unlikely that nickel in house dust results in a potential health risk.

**Zinc.** The background exposure to zinc via food intake accounts for 70% of the TDI, whereas exposure via house dust usually represents only a minor exposure route (2.1% of the TDI for children is filled up for the P95-concentration in house dust). The risk index was only exceeded for the maximum concentration, which seems to be an exceptionally high concentration with regard to other data and concentrations determined in house dust in the Netherlands. This indicates that it is highly unlikely that zinc in house dust results in a potential health risk.

**Di(2-ethylhexyl)-phthalate (DEHP).** The background exposure to DEHP due to consumption of food and drinking water accounts for 24-52% of the TDI for a child and 6-32% for an adult. The extra exposure via house dust may lead to marked exceedance of the TDI for both adults and especially children. Given the toxicological basis for the TDI for DEHP with disruption of reproductive development in males as the critical effect, lower levels of exposure later in life do not compensate for exceedance of the TDI during child age. Accordingly exposure should not be integrated over an entire life for DEHP when comparing to its TDI. In conclusion, DEHP poses a human health risk during child age, both due to the background exposure and due to house dust intake.

**Sum PAHs.** In the present report the so-called indicator approach was used for PAHs, which involves using benzo(a)pyrene as the marker for the total mixture, both in quantifying risk and exposure. Background exposure to PAHs is low compared to the Maximum Permisible Risk (MPR) (12% of the MPR is filled up by the background exposure for an adult and 30% for a child). Remarkably the MPR is sometimes exceeded due to exposure via house dust. Given the general policy of making exposures to the genotoxic and carcinogenic PAH via food as low as reasonably possible, this potentially high exposure via house dust is significant.

**BDE99.** A potential health risk due to exposure via house dust is identified for the only congener for which a toxicological reference value is available (BDE99). This is shown by the risk indices for adults and children respectively of 0.45 and 1.5 for the geometric mean concentration data in house dust. The contribution of house dust to the exposure to BDE99 may be considerable, especially for children. For example, 6% and 57% of the allowable daily intake level may be filled up by dust intake for adults and children based on the EU geometric mean data. For comparison of the exposure to the provisional toxicological reference dose, the exposure over an entire lifetime should be considered. As the expected exposure due to dust (and food) intake is expected to exceed the maximum allowable intake level on a regular basis both for adults and children, a potential health risk due to BDE99 is expected to occur regularly.

Comparison between concentrations of BDE209 to other BDE-congeners in house dust indicates that human exposure to BDE209 via house dust may be (much) greater than to the other congeners, and is probably also much greater than exposure to BDE209 via food intake. In order to arrive at more definitive conclusions a toxicological reference value for

BDE209 in particular should become available, which is also highly recommended for the other PBDE congeners. In order to derive toxicological reference values additional toxicity studies are required.

Measures to decrease human exposure to substances in house dust. Little is known about measures to prevent human exposure to substances via house dust. If possible the source of the substances in house dust should be identified and measures should be taken to prevent further contamination in the future. Dust control may be effective if performed properly (preferably by wet cleaning) and regularly. The measures should be evaluated after some time by analysis of dust samples as rebouncing to former substance concentrations may occur.

### Acknowledgement

The authors would like to thank Cor van den Bogaard (Inspectorate of the Ministry of Housing, Spatial Planning and the Environment of the Netherlands), Annelike Dusseldorp, Marco Zeilmaker, Bert-Jan Baars, and Adriënne Sips (all RIVM) for reviewing the report.

### References

- Adgate JL, Rhoads GG, Lioy PJ (1998a) The use of isotope ratios to apportion sources of lead in Jersey City, NJ, house dust wipe samples. Sci Total Environ 221, 171-180.
- Adgate JL, Willis RD, Buckley TG, Chow JC, Watson J, Rhoads GG, Lioy PJ (1998b) Chemical mass balance source apportionment of lead in house dust. Environ Sci Technol 32, 108-114.
- Al Bitar F (2004) Hazardous chemicals in Belgian house dust. www.greenpeace.org/raw/content/belgium/fr/press/reports/hazardous-chemicals-in-belgium-2.pdf. Retrieved on 19-09-2007
- ATSDR (1999) Toxicological Profile for Uranium. US Agency of Toxic Substances and Disease Registry.
- Australian Government (1992) ADI list: Acceptable Daily Intakes for Agricultural and Veterinary Chemicals. http://www.tga.gov.au/docs/pdf/adi.pdf. Retrieved on 18-12-2007.
- Baars AJ, Theelen RMC, Janssen PJCM, Hesse JM, Apeldoorn ME van, Meijerink MCM, Verdam L, Zeilmaker MJ (2001) Re-evaluation of human-toxicological Maximum Permissible Levels. Report no. 711701025, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Bakker MI, De Winter-Sorkina R, De Mul A, Boon PE, Van Donkersgoed G, Van Klaveren JD, Baumann BA, Hijman WC, Van Leeuwen SPJ, De Boer J, Zeilmaker MJ (2008) Dietary intake and risk evaluation of polybrominated diphenyl ethers in The Netherlands. Mol. Nutr. Food Res 52, 204-216.
- Beamer P, Castano A, Leckie JO (2002) Vertical profile particulate matter measurements in a California daycare. Proceedings: Indoor Air 2002, 103-108.
- Becker K, Seiwert M, Angerer J, Heger W, Koch HM, Nagorka R, Rosskamp E, Schluter C, Seifert B, Ullrich D (2004) DEHP metabolites in urine of children and DEHP in house dust. International Journal of Hygene and Environmental Health 207, 409-417.
- Becker K, Seiwert M, Angerer J, Kolossa-Gehring M, Hoppe HW, Ball M, Schulz C, Thumulla J, Seifert B (2006) GerES IV pilot study: assessment of the exposure of German children to organophosphorus and pyrethroid pesticides. International Journal of Hygene and Environmental Health 209, 221-233.
- Berger-Preiß E, Levsen K, Leng G, Idel H, Sugiri D, Ranft U (2002) Indoor pyrethroid exposure in homes with woollen textile floor coverings. International Journal of Hygene and Environmental Health 205, 459-472.
- BfR (2006) Pflanzenschutzmittel-Wirkstoffe: ADI-Werte und gesundheitliche Trinkwasser-Leitwerte Aktualisierte Information. http://www.bfr.bund.de/cm/218/pflanzenschutzmittel wirkstoffe adi werte und gesundheitli

- che trinkwasser leitwerte.pdf. Retrieved on 18-12-2007.
- Bornehag CG, Lundgren B, Weschler CJ, Sigsgaard T, Hagerhed-Engman L, Sundell J (2005)

  Phthalates in indoor dust and their association with building characteristics. Environ Health Perspect 113, 1399-1404.
- Bornehag CG, Sundell J, Weschler CJ, Sigsgaard T, Lundgren B, Hasselgren M, Hagerhed-Engman L (2004) The association between asthma and allergic symptoms in children and phthalates in house dust: a nested case-control study. Environ Health Perspect 112, 1393-1397.
- Bruggen M van, Janssen PCJM, Kliest JJG, Meulenbelt J, Smetsers RCGM, Uijt de Haag PAM, Mik G de (1998) Gezondheidsrisico's brand El Al-Boeing. Report no. 609026002, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Butte W, Heinzow B (2002) Pollutants in house dust as indicators of indoor contamination. Rev Environ Contam Toxicol 175, 1-46.
- Butte W, Hoffmann W, Hostrup O, Schmidt A, Walker G (2001) Endokrin wirksame Substanzen im Hausstaub: Ergebnisse eines repräsentativen Monitorings. Gefahrst. Reinhalt. Luft 61, 19-23.
- Calabrese EJ, Barnes R, Stanek EJ 3rd, Pastides H, Gilbert CE, Veneman P, Wang XR, Lasztity A, Kostecki PT (1989) How much soil do young children ingest: an epidemiologic study. Regul Toxicol Pharmacol 10, 123-137.
- Campbell C, Schwarz DF, Rich D, Dockery DW (2003) Effect of a follow-up professional home cleaning on serial dust and blood lead levels of urban children. Arch Environ Health 58, 771-780.
- Carrizales L, Razo I, Téllez-Hernández JI, Torres-Nerio R, Torres A, Batres LE, Cubillas A-C, Díaz-Barriga F (2005) Exposure to arsenic and lead of children living near a copper-smelter in San Luis Potosi, Mexico: Importance of soil contamination for exposure of children. Environmental Research 101, 1-10.
- Chuang JC, Callahan PJ, Lyu CW, Wilson NK (1999) Polycyclic aromatic hydrocarbon exposures of children in low-income families. J Expo Anal Environ Epidemiol 9, 85-98.
- Clausen PA, Lindeberg Bille RL, Nilsson T, Hansen V, Svensmark B, Bowadt S (2003) Simultaneous extraction of di(2-ethylhexyl) phthalate and nonionic surfactants from house dust.

  Concentrations in floor dust from 15 Danish schools. J Chromatogr A 986, 179-190.
- Costner P, Thorpe B, McPherson A (2005) Sick of dust: Chemicals in common products A needless health risk in our homes. www.cleanproduction.org/library/Dust%20Report%20with%20 Appendices.pdf. Retrieved on 11-09-2007.
- Curwin BD, Hein MJ, Sanderson WT, Nishioka MG, Reynolds SJ, Ward EM, Alavanja MC (2005) Pesticide contamination inside farm and nonfarm homes. J Occup Environ Hyg 2, 357-367.
- Davis JJ, Gulson BL (2005) Ceiling (attic) dust: a "museum" of contamination and potential hazard. Environ Res 99, 177-194.

- De Boer E (2007) Student's report at the RIVM-SIR and VU: House dust: composition, estimation of human exposure, available measurement techniques and the measurement of bioavailability of dust related Poly Brominated Fire Retardants.
- De Boer J, Wells DE (2006) Pitfalls in the analysis of brominated flame retardants in environmental, human and food samples including results of three international interlaboratory studies. Trends in Analytical Chemistry 25, 364-372.
- De Winter-Sorkina R, Bakker MI, Wolterink G, Zeilmaker MJ (2006) Brominated flame retardants: occurence, dietary intake and risk assessment. Report no. 320100002, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, the Netherlands.
- Dutch Health Council (Gezondheidsraad) (1997) Committee on lead in drinking water. Report no. 1997/07.
- EFSA (2005) Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids and Materials in Contact with Food (AFC) on a request from the Commission related to Bis(2-ethylhexyl)phthalate (DEHP) for use in food contact materials Question N° EFSA-Q-2003-191. The EFSA Journal 243, 1-20.
- EFSA (2005a) Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids and Materials in Contact with Food (AFC) related to Butylbenzylphthalate (BBP) for use in food contact materials. The EFSA Journal 241: 1-14.
- EFSA (2005b) Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids and materials in Contact with Food (AFC) related to Bis(2-ethylhexyl)phthalate (DEHP) for use in food contact materials. The EFSA Journal 243: 1-20.
- EFSA (2005c) Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to Di-isodecylphthalate (DIDP) for use in food contact materials. The EFSA Journal 245: 1-14.
- EFSA (2005d) Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to Di-isononylphthalate (DINP) for use in food contact materials. The EFSA Journal 244: 1-18.
- EU (2003) Review report for the active substance pendimethalin. European Commission Health & Consumer Protection Directorate-General. Pendimethalin 7477/VI/98-final, dated 13 January 2003. http://ec.europa.eu/food/plant/protection/evaluation/existactive/list1-35\_en.pdf. Retrieved on 18-12-2007.
- EU (2006) Review report for the active substance phosmet. Finalised in the Standing Committee on the Food Chain and Animal Health at its meeting on 24 November 2006 in view of the inclusion of phosmet in Annex I of Directive 91/414/EEC. http://ec.europa.eu/food/plant/protection/evaluation/existactive/list\_phosmet.pdf. Retrieved on 18-12-2007.
- Fenske RA, Lu C, Barr D, Needham L (2002) Children's exposure to chlorpyrifos and parathion in an agricultural community in central Washington State. Environ Health Perspect 110, 549-553.

- Fergusson JE, Kim ND (1991) Trace elements in street and house dust: sources and speciation. The Science of the Total Environment 100, 125-150.
- Fergusson JE, Schroeder RJ (1985) Lead in house dust of Christchurch, New Zealand: sampling, levels and sources. Sci Total Environ 46, 61-72.
- Food Standards Agency. www.food.gov.uk/multimedia/pdfs/evm\_magnesium.pdf. Retrieved on 18-12-2007.
- Freeman NC, Stern AH, Lioy PJ (1997) Exposure to chromium dust from homes in a Chromium Surveillance Project. Arch Environ Health 52, 213-219.
- Fromme H, Lahrz T, Piloty M, Gebhart H, Oddoy A, Ruden H (2004) Occurrence of phthalates and musk fragrances in indoor air and dust from apartments and kindergartens in Berlin (Germany). Indoor Air 14, 188-195.
- Fromme H, Mattulat A, Lahrz T, Ruden H (2005) Occurrence of organotin compounds in house dust in Berlin (Germany). Chemosphere 58, 1377-1383.
- Gevao B, Al-Bahloul M, Al-Ghadban AN, Al-Omair A, Ali L, Zafar J, Helaleh M (2006) House dust as a source of human exposure to polybrominated diphenyl ethers in Kuwait. Chemosphere 64, 603-608.
- Gezondheidsraad (Dutch Health Council) (1997), Lead in drinking water. Report no. 1997/07.
- Gulson BL, Davis JJ, Mizon KJ, Korsch MJB-SJ (1995) Sources of lead in soil and dust and the use of dust fallout as a sampling medium. The Science of the Total Environment 166, 245-262.
- Harrad S, Hazrati S, Ibarra C (2006) Concentrations of polychlorinated biphenyls in indoor air and polybrominated diphenyl ethers in indoor air and dust in Birmingham, United Kingdom: implications for human exposure. Environ Sci Technol 40, 4633-4638.
- Harrad S, Ibarra C, Diamond M, Melymuk L, Robson M, Douwes J, Roosens L, Dirtu AC, Covaci A (2008) Polybrominated diphenyl ethers in domestic indoor dust from Canada, New Zealand, United Kingdom and United States. Environment International 34, 232-238.
- Hawley JK (1985) Assessment of health risk from exposure to contaminated soil. Risk Analysis 5, 289-302.
- Hoeven PH and Engelen JGM van RIVM (1997) Dicamba- Residue evaluation and risk assessment for the consumer. RIVM/CSR Advisory report: 05229A00, August 1997 (confidential).
- Hogenkamp A, Vaal M, Heederik D (2004) Pesticide exposure in dwellings near bulb growing fields in The Netherlands: an explorative study. Ann Agric Environ Med 11, 149-153.
- Hysong TA, Burgess JL, Cebrian Garcia ME, O'Rourke MK (2003) House dust and inorganic urinary arsenic in two Arizona mining towns. J Expo Anal Environ Epidemiol 13, 211-218.
- Ibarra C, Hazrati S, Harrad S (2006) Polybrominated Diphenyl Ethers (PBDEs) and Polychlorinated Biphenyls (PCBs) in household dust in Birmingham, UK. Poster presentation, Dioxin meeting,

64

- august 2006.
- IPCS (International Programme on Chemical Safety) (2001) Arsenic and Arsenic Compounds (Second Edition); Environmental Health Criteria 224. World Health Organization, Geneva, Switzerland
- Janssen PJCM, Apeldoorn ME van, Koten-Vermeulen JEM van, Mennes WC (1995) Human-Toxicological Criteria for Serious Soil Contamination: Compounds evaluated in 1993 & 1994. Report no. 320007001, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Janssen PJCM, Apeldoorn ME van, Engelen JGM van, Schielen PCJI, Wouters MFA (1998) Maximum Permissible Risk levels for human intake of soil contaminants: Fourth series of compounds. Report no. 711701004, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Janssen PJCM and Baars AJ (2004) Oriënterende Evaluatie Gezondheidsrisico Metalen in Tatoeages. Report no. 320105001, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- JMPR (1991) http://www.who.int/ipcs/publications/jmpr/pesticide\_inventory.pdf. Retrieved on 18-12-2007.
- JMPR (1995) http://www.who.int/ipcs/publications/jmpr/pesticide\_inventory.pdf. Retrieved on 17-12-2007.
- JMPR (2001) http://www.inchem.org/documents/jmpr/jmpeval/jmpr2002.htm. Retrieved on 17-12-2007.
- JMPR (2002) http://www.who.int/ipcs/publications/jmpr/pesticide\_inventory.pdf. Retrieved on 17-12-2007.
- JMPR (2003) http://www.who.int/ipcs/publications/jmpr/pesticide\_inventory.pdf. Retrieved on 17-12-2007.
- JMPR (2004a) http://www.fao.org/ag/AGP/AGPP/Pesticid/JMPR/DOWNLOAD/2004\_rep/report2004jmpr.pdf. Retrieved on 17-12-2007.
- JMPR (2004b) http://www.who.int/ipcs/publications/jmpr/pesticide\_inventory.pdf. Retrieved on 17-12-2007.
- JMPR (2005) http://www.fao.org/docrep/009/a0209e/a0209e0a.htm. Retrieved on 17-12-2007.
- JMPR (2006) http://www.who.int/ipcs/publications/jmpr/pesticide\_inventory.pdf. Retrieved on 17-12-2007.
- Jones-Otazo HA, Clarke JP, Diamond ML, Archbold JA, Ferguson G, Harner T, Richardson GM, Ryan JJ, Wilford B (2005) Is house dust the missing exposure pathway for PBDEs? An analysis of the urban fate and human exposure to PBDEs. Environ Sci Technol 39, 5121-5130.

- Jovanovic S, Carrot F, Deschamps C, Deschamps N, Vukotic P (1995) A study of the air-pollution in the surroundings of an aluminium smelter, using epiphytic and lithophytic lichens. J Trace Microprobe Tech 13, 463-471.
- Knoth W, Mann W, Meyer R, Nebhuth J (2003) Brominated diphenyl ether in indoor dust. Organohalogen Compounds 61, 207-210.
- Lanphear BP, Succop P, Roda S, Henningsen G (2003) The effect of soil abatement on blood lead levels in children living near a former smelting and milling operation. Public Health Rep 118, 83-91.
- Lapitajs G, Greb U, Dunemann L, Begerow J, Moens L, Verrept P (1995). ICPMS in the determination of trace and ultratrace elements in the human body. Int Lab 25, 21–27.
- Lewis RG, Fortmann RC, Camann DE (1994) Evaluation of methods for monitoring the potential exposure of small children to pesticides in the residential environment. Arch Environ Contam Toxicol 26, 37-46.
- Lewis RG, Fortune CR, Blanchard FT, Camann DE (2001) Movement and deposition of two organophosphorus pesticides within a residence after interior and exterior applications. J Air Waste Manag Assoc 51, 339-351.
- Lewis RG, Fortune CR, Willis RD, Camann DE, Antley JT (1999) Distribution of pesticides and polycyclic aromatic hydrocarbons in house dust as a function of particle size. Environ Health Perspect 107, 721-726.
- Lijzen JPA, Baars AJ, Otte PF, Rikken MGJ, Swartjes FA, Verbruggen EMJ, van Wezel AP (2001) Technical evaluation of the intervention values for soil/sediment and groundwater. Human and ecotoxicological risk assessment and derivation of risk limits for soil, aquatic sediment and groundwater. Report no. 711701023, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Lu C, Fenske RA, Simcox NJ, Kalman D (2000) Pesticide exposure of children in an agricultural community: evidence of household proximity to farmland and take home exposure pathways. Environ Res 84, 290-302.
- Lu C, Kedan G, Fisker-Andersen J, Kissel JC, Fenske RA (2004) Multipathway organophosphorus pesticide exposures of preschool children living in agricultural and nonagricultural communities. Environ Res 96, 283-289.
- Maertens RM, Bailey J, White PA (2004) The mutagenic hazards of settled house dust: a review. Mutat Res 567, 401-425.
- Meyer I, Heinrich J, Lippold U (1999) Factors affecting lead, cadmium, and arsenic levels in house dust in a smelter town in Eastern Germany. Environ. Res. 81, 32-44.
- Mushak P (1998) Uses and limits of empirical data in measuring and modeling human lead exposure. Environ. Health Perspect. 106, 1467-1484.
- Obendorf SK, Lemley AT, Hedge A, Kline AA, Tan K, Dokuchayeva T (2006) Distribution of

- pesticide residues within homes in central New York State. Arch Environ Contam Toxicol 50, 31-44.
- Oie L, Hersoug LG, Madsen JO (1997) Residential exposure to plasticizers and its possible role in the pathogenesis of asthma. Environ Health Perspect 105, 972-978.
- Oomen AG, Janssen PJCM, Van Eijkeren JCH, Bakker MI, Baars AJ (2007) Cadmium in de Kempen: een integrale risicobeoordeling. Report no. 320007001, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Oomen AG, Lijzen JPA (2004) Relevancy of human exposure via house dust to the contaminants lead and asbestos. Report no: 711701037, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Otte PF, Lijzen JPA, Otte JG, Swartjes FA, Versluijs CW (2001) Evaluation and revision of the CSOIL parameter set. Report no. 711701021, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Paustenbach DJ, Finley BL, Long TF (1997) The critical role of house dust in understanding the hazards posed by contaminated soils. International Journal of Toxicology 16, 339-362.
- Pless-Mulloli T, Schecter A, Schilling B, Paepke O (2006) Levels of PBDE in household dust and lint in the UK, Germany and the USA. Poster presentation, Dioxin meeting, august 2006.
- Pohner A, Simrock S, Thumulla J, Weber S, Wirkner T (1997) Hintergrundbelastung des hausstaubes von privathauhalten mit mittel- und schwerfluchtigen organischen schadstoffen (in German). Umwelt Gesundheit 2, 1-64.
- Rasmussen PE, Subramanian KS, Jessiman BJ (2001) A multi-element profile of housedust in relation to exterior dust and soils in the city of Ottawa, Canada. Sci Total Environ 267, 125-140.
- Rasmussen PE, Wheeler AJ, Hassan NM, Filiatreault A, Lanouette M (2007) Monitoring personal, indoor, and outdoor exposures to metals in airborne particulate matter: risk of contamination during sampling, handling and analysis. Atmospheric Environment 41, 5897-5907.
- Regueiro J, Llompart M, Garcia-Jares C, Cela R (2006) Determination of polybrominated diphenyl ethers in domestic dust by microwave-assisted solvent extraction and gas chromatographytandem mass spectrometry. J Chromatogr A 1137, 1-7.
- Roberts JW, Dickey P (1995) Exposure of children to pollutants in house dust and indoor air. Rev Environ Contam Toxicol 143, 59-78.
- Rothlein J, Rohlman D, Lasarev M, Phillips J, Muniz J, McCauley L (2006) Organophosphate pesticide exposure and neurobehavioral performance in agricultural and non-agricultural Hispanic workers. Environ Health Perspect 114, 691-696.
- Santillo D, Labunska I, Davidson H, Johnson P, Strutt M, Knowles O (2003) Consuming chemicals: Hazardous chemicals in house dust as an indicator of chemical exposure in the home.
- SCF (2002) Opinion of the Scientific Committee on Food on the risks to human health of Polycyclic

- Aromatic Hydrocarbons in food. SCF/CS/CNTM/PAH/29.
- Schecter A, Papke O, Joseph JE, Tung KC (2005) Polybrominated diphenyl ethers (PBDEs) in U.S. computers and domestic carpet vacuuming: possible sources of human exposure. J Toxicol Environ Health A 68, 501-513.
- Seifert B, Becker K, Helm D, Krause C, Schulz C, Seiwert M (2000) The German Environmental Survey 1990/1992 (GerES II): reference concentrations of selected environmental pollutants in blood, urine, hair, house dust, drinking water and indoor air. J Expo Anal Environ Epidemiol 10, 552-565.
- Sharp R, Lunder S (2004) In the dust: toxic fire retardants in American homes. http://www.ewg.org/reports/inthedust/index.php. Retrieved on 11-09-2007.
- Simcox NJ, Fenske RA, Wolz SA, Lee IC, Kalman DA (1995) Pesticides in household dust and soil: exposure pathways for children of agricultural families. Environ Health Perspect 103, 1126-1134.
- Sjodin A, Papke O, Focant J-F, Jones RS, Pless-Mulloli T, Toms L-M, Wang R, Zhang Y, Herrmann T, Patterson DG Jr (2006) Concentration of polybrominated diphenyl ethers (PBDEs) in household dust from various countries is dust a major source of human exposure? Poster presentation, Dioxin meeting, august 2006.
- Sjodin A, Papke O, McGahee E, Jones R, Focant J-F, Pless-Mulloli T, Toms L-M, Wang R, Zhang Y, Needham L, Herrmann T, Patterson DG (2004) Concentration of polybrominated diphenyl ethers (PBDEs) in household dust from various countries Inhalation a potential route of human exposure. Organohalogen Compounds 66, 3818-3822.
- Spalinger SM, von Braun MC, Petrosyan V, von Lindern IH (2007) Northern Idaho house dust and soil lead levels compared to the Bunker Hill Superfund Site. Environ Monit Assess 130, 57-72.
- Stanek EJ, Calabrese EJ (1995) Soil ingestion estimates for use in site evaluations based on the best tracer method. Human and Ecological Risk Assessment 1, 133-156.
- Stapleton HM, Dodder NG, Offenberg JH, Schantz MM, Wise SA (2005) Polybrominated diphenyl ethers in house dust and clothes dryer lint. Environ Sci Technol 39, 925-931.
- Sterling DA, Roegner KC, Lewis RD, Luke DA, Wilder LC, Burchette SM (1999) Evaluation of four sampling methods for determining exposure of children to lead-contaminated household dust. Environmental Research Section A 81, 130-141.
- Tan J, Cheng SM, Loganath A, Chong YS, Obbard JP (2007a) Polybrominated diphenyl ethers in house dust in Singapore. Chemosphere 66, 985-992.
- Tan J, Cheng SM, Loganath A, Chong YS, Obbard JP (2007b) Selected organochlorine pesticide and polychlorinated biphenyl residues in house dust in Singapore. Chemosphere 68, 1675-1682.
- Thatcher TL, Layton DW (1995) Deposition, resuspension, and penetration of particles within a residence. Atmospheric Environment 29, 1487-1497.

- Toms L, Muller J, Bartkow M, Symons R (2006) Brominated flame retardants in indoor environments in Australia.
- Van Engelen JGM, Park MVDZ, Janssen PJCM, Oomen AG, Brandon EFA, Bouma K, Sips AJAM, Van Raaij MTM (in press) Chemicals in Toys: A general methodology for assessment of chemical safety of toys with a focus on elements. Report no. 320003001.
- USEPA (1989) IRIS-file 1,1-biphenyl. http://www.epa.gov/ncea/iris/subst/0013.htm. Retrieved on 18-12-2007.
- USEPA (1993) IRIS-file Atrazin. http://www.epa.gov/NCEA/iris/subst/0209.htm. Retrieved on 17-12-2007.
- USEPA (1998) Reregistration Eligibility Decision (RED) Alachlor. http://epa.gov/oppsrrd1/REDs/0063.pdf. Retrieved on 17-12-2007.
- USEPA (1997) National Center for Environmental Assessment: Washington DC.
- USEPA (2002) National Center for Environmental Assessment: Washington DC.
- WHO (1996) Drinking-water Guidelines- 2<sup>nd</sup> edition. http://www.who.int/water sanitation health/dwq/2edvol2p2d.pdf. Retrieved on 17-12-2007.
- WHO (2003) Metolachlor in Drinking-water. Background document for development of WHO Guidelines for Drinking-water Quality. http://www.who.int/water\_sanitation\_health/dwq/chemicals/metolachlor.pdf. Retrieved on 17-12-2007.
- Wilford BH, Shoeib M, Harner T, Zhu J, Jones KC (2005) Polybrominated diphenyl ethers in indoor dust in Ottawa, Canada: implications for sources and exposure. Environ Sci Technol 39, 7027-7035.
- Williams S (2002) ATSDR, Health Consultation: Review of Dust Sampling Data for 45th Street Artists' Cooperative. http://www.atsdr.cdc.gov/hac/PHA/artistscoop/wil\_p2.html. Retrieved on 20-12-2007.
- Wilson NK, Chuang JC, Lyu C, Menton R, Morgan MK (2003) Aggregate exposures of nine preschool children to persistent organic pollutants at day care and at home. J Expo Anal Environ Epidemiol 13, 187-202.
- Winter-Sorkina R de, Bakker MI, Wolterink G, Zeijlmaker MJ (2006) Brominated flame retardants: occurence, dietary intake and risk assessment. Report no. 320100002, available at http://www.rivm.nl/en/, National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Wormuth M, Scheringer M, Vollenweider M, Hungerbuhler K (2006) What are the sources of exposure to eight frequently used phthalic acid esters in Europeans? Risk Anal 26, 803-824.
- Yiin LM, Rhoads GG, Rich DQ, Zhang J, Bai Z, Adgate JL, Ashley PJ, Lioy PJ (2002) Comparison of techniques to reduce residential lead dust on carpet and upholstery: the new jersey assessment of cleaning techniques trial. Environ Health Perspect 110, 1233-1237.

Zhu J, Feng Y-L, Shoeib M (2007) Detection of Dechlorane plus in residential indoor dust in the city of Ottawa, Canada. Environ. Sci Technol 41, 7694-7698.

### **Appendix 1. References on substances in house dust**

Table Appendix 1. Information on the references in which information on the concentrations of substances in house dust was found.

Substance	Concentration in dust (mg/kg)			Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Metals							
Aluminium	24281	51100	44225	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Antimony	25.5	66.4		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Antimony	5.5	14.7		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Antimony	5.54	57.4	15.4	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Arsenic	2.6	57.1		container	Germany-Hettstedt	(Meyer et al., 1999)	
Arsenic	9.00			vacuum	USA-Utah	(Lanphear <i>et al.</i> , 2003)	
Arsenic	26.3	79.7		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Arsenic	12.2	17.6		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Arsenic	70.1	192	114.2 (P90)	vacuum	USA-Hayden	(Hysong et al., 2003)	Copper mining/smelting town
Arsenic	46.7	130	88 (P90)	vacuum	USA-Winkelman	(Hysong et al., 2003)	Copper mining/smelting town
Arsenic	4.9	79.5	18.5	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Arsenic	2.09	182	6.7	vacuum	Germany	(Seifert et al., 2000)	
Barium	454	1480	803	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Beryllium	0.53	1.0	0.90	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Bismuth	1.02	8.62	6.48	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Cadmium	2.6	52.7		container	Germany-Hettstedt	(Meyer et al., 1999)	
Cadmium	13	52.0		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting

Substance	Concer	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric						
	mean	Maximum	P95				
Metals							
Cadmium	2.2	3.0		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Cadmium	4.42	34.9	17.3	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Cadmium	0.86	220	5.6	vacuum	Germany	(Seifert et al., 2000)	
Chromium	111	188		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Chromium	159	5440		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Chromium	66			wipe	USA-New Jersey	(Freeman <i>et al.</i> , 1997)	
Chromium	75.4	330.3	191.8	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Chromium	64.2	1330	178	vacuum	Germany	(Seifert et al., 2000)	
Cobalt	17	20		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Cobalt	12	20		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Cobalt	8.40	22.7	13.1	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Copper	261	396		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Copper	120	364		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Copper	171	601	489	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Copper	79.9	12540	339	vacuum	Germany	(Seifert et al., 2000)	
Lead	1200	18600		wipe	USA-New Jersey	(Adgate et al., 1998a)	
Lead	857			vacuum		(Adgate et al., 1998b)	
Lead	1660	2594		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Lead	477	1150		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Lead	127			vacuum	USA-Utah	(Lanphear <i>et al.</i> , 2003)	
Lead	128	1947		container	Germany-Hettstedt	(Meyer et al., 1999)	
Lead	233	3226	1312	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Lead	5.9	37000	178	vacuum	Germany	(Seifert et al., 2000)	

Substance	Concer	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Metals							
Lead	120	830		vacuum	USA-Idaho	(Spalinger <i>et al.</i> , 2007)	Average of all towns measured
Lithium	6.1	15.5	8.2	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Magnesium	9442	23250	13390	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Magnesium	2110	52000	6400	vacuum	Germany	(Seifert et al., 2000)	
Manganese	260.3	423.5	407.3	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Manganese	109	9410	333	vacuum	Germany	(Seifert et al., 2000)	
Mercury	1.73	37,099	12,558	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Molybdenum	2.8	6.5		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Molybdenum	1.5	6.7		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Molybdenum	1.96	28.64	14.22	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Nickel	47	80		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Nickel	26	50		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Nickel	53.6	243.3	116.4	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Rubidium	24.6	40.2	34.9	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Selenium	1.0	6.8	2.2	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Silver	1.48	9.33	6.50	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	

Substance	Concer	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Metals							
Strontium						(Rasmussen et al.,	
	242	410	382	vacuum	Canada-Ottawa	2001)	
Strontium	32.1	1170	119	vacuum	Germany	(Seifert et al., 2000)	
						(Rasmussen et al.,	
Tellurium	0.07	0.28	0.13	vacuum	Canada-Ottawa	2001)	
						(Rasmussen et al.,	
Thallium	0.14	0.24	0.21	vacuum	Canada-Ottawa	2001)	
						(Rasmussen et al.,	
Tin	21.87	595.02	221.33	vacuum	Canada-Ottawa	2001)	
Titanium	2854	4527		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Titanium	2835	4983		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Tungsten	3.7	5.6		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Tungsten	2.8	5.2		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Uranium	0.55	1.33	1.06	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Vanadium	112	163		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Vanadium	107	193		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Vanadium	23.7	43.6	39.9	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Zinc	17294	57400		ceiling brush	Australia	(Davis et al., 2005)	Industrial setting
Zinc	577	1160		ceiling brush	Australia	(Davis et al., 2005)	Non-industrial setting
Zinc	628.0	1840	1460.8	vacuum	Canada-Ottawa	(Rasmussen <i>et al.</i> , 2001)	
Zinc	475	30600	1570	vacuum	Germany	(Seifert et al., 2000)	
Organotin compounds							
Dibutyltin (DBT)	0.51	5.60	3.28	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2005)	
Dioctyltin (DOT)	0.02	0.36	0.03	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2005)	
Monobutyltin (MBT)	0.16	1.50	0.70	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2005)	

Substance	Concer	ntration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Organotin compounds							
Monooctyltin (MOT)	0.01	0.04	0.03	vacuum	Germany-Berlin	(Fromme et al., 2005)	
Tributyltin (TBT)	0.02	0.08	0.07	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2005)	
Pesticides							
2,4-D	1.24	7.29		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
2,4-D	330 μg/kg			surface sampler	USA-Iowa	(Curwin et al., 2005)	Non-farm
2,4-D		1.5		vacuum	USA-New York state	(Obendorf <i>et al.</i> , 2006)	
Alachlor		1.5		vacuum	USA-New York state	(Obendorf et al., 2006)	
Aldrin	0.006	0.051		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
alpha-Chlordane <sup>1</sup>	0.055	0.256		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
alpha-HCH	0.7 μg/kg	0.0087		airconditioning	Singapore	(Tan et al., 2007b)	
Altazine		0.5		vacuum	USA-New York state	(Obendorf et al., 2006)	
Atrazine	2.3 μg/kg			surface sampler	USA-Iowa	(Curwin et al., 2005)	Non-farm
Azinphos methyl	0.29	1.1		vacuum	USA-Washington	(Lu et al., 2000)	Non-farm
Azinphos methyl	0.330	0.816		surface sampler	USA-Washington	(Simcox et al., 1995)	Non-farm
AZM	5.9	16		surface sampler	USA-Oregon	(Rothlein <i>et al.</i> , 2006)	
beta-HCH	2.2 μg/kg	0.057		airconditioning	Singapore	(Tan et al., 2007b)	
Carbaryl		1.0		vacuum	USA-New York state	(Obendorf et al., 2006)	
Chloroprofam		0.17		vacuum	The Netherlands	(Hogenkamp <i>et al.</i> , 2004)	Non-farm
Chlorpyrifos	1.04	6.45		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Chlorpyrifos	0.54			vacuum	USA-Washington	(Fenske et al., 2002)	

Substance	Concer	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric		· · · · · ·				
	mean	Maximum	P95				
Pesticides							
Chlorpyrifos	0.38			vacuum	USA-Washington	(Fenske et al., 2002)	
Chlorpyrifos	30 μg/kg			surface sampler	USA-Iowa	(Curwin et al., 2005)	Non-farm
Chlorpyrifos		1.7		vacuum	USA-New York state	(Obendorf <i>et al.</i> , 2006)	
Chlorpyrifos	0.20	1.2		surface sampler	USA-Oregon	(Rothlein <i>et al.</i> , 2006)	
Chlorpyrifos	0.168	0.483		surface sampler	USA-Washington	(Simcox et al., 1995)	Non-farm
Cis- Chlordane <sup>1</sup>	2.6 μg/kg	0.039		airconditioning	Singapore	(Tan et al., 2007b)	
DDD	4.7 μg/kg	0.048		airconditioning	Singapore	(Tan et al., 2007b)	
DDE	0.007	0.047		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
DDE	5.9 μg/kg	0.05		airconditioning	Singapore	(Tan et al., 2007b)	
DDT	0.121	0.782		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
DDT	56 μg/kg	0.7		airconditioning	Singapore	(Tan et al., 2007b)	
delta-HCH	5.5 μg/kg	0.17		airconditioning	Singapore	(Tan et al., 2007b)	
Diazinon	0.044	0.216		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Diazinon	0.14	0.61		surface sampler	USA-Seattle- Washington	(Lu et al., 2004)	Non-farm
Diazinon		2.0		vacuum	USA-New York state	(Obendorf et al., 2006)	
Diazinon	0.31	0.72		surface sampler	USA-Oregon	(Rothlein et al., 2006)	
Dicamba		2.5		vacuum	USA-New York state	(Obendorf et al., 2006)	
Dieldrin	0.018	0.050		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Dimethyl OP	0.37	1.3		vacuum	USA-Washington	(Lu et al., 2000)	Non-farm
Ethyl parathion	0.076	0.425		surface sampler	USA-Washington	(Simcox et al., 1995)	Non-farm
Ethyl parathion	0.56			vacuum	USA-Washington	(Fenske et al., 2002)	
Ethyl parathion	0.04			vacuum	USA-Washington	(Fenske et al., 2002)	
gamma-Chlordane <sup>2</sup>	0.098	0.471		surface sampler	USA-North Carolina	(Wilson et al., 2003)	

Substance	Concen	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Pesticides							
gamma-HCH <sup>3</sup>	2.9 μg/kg	0.074		airconditioning	Singapore	(Tan et al., 2007b)	
Glyphosate	140 μg/kg			surface sampler	USA-Iowa	(Curwin et al., 2005)	Non-farm
Heptachlor	0.119	0.335		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Lindane <sup>3</sup>	0.33	0.046		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Malathion		2.0		vacuum	USA-New York state	(Obendorf et al., 2006)	
Malathion	0.38	1.4		surface sampler	USA-Oregon	(Rothlein et al., 2006)	
Mecoprop		0.4		vacuum	USA-New York state	(Obendorf et al., 2006)	
Methamidophos		0.4		vacuum	USA-New York state	(Obendorf et al., 2006)	
Methyl parathion		0.125		vacuum	USA-New York state	(Obendorf et al., 2006)	
Methyl parathion	0.38	1.9		surface sampler	USA-Oregon	(Rothlein et al., 2006)	
Metolachlor	5.7 μg/kg			surface sampler	USA-Iowa	(Curwin et al., 2005)	Non-farm
Metolachlor		0.8		vacuum	USA-New York state	(Obendorf et al., 2006)	
Pendimethalin		3.0		vacuum	USA-New York state	(Obendorf et al., 2006)	
Permethrin	0.14	187.00	11.5	vacuum	Germany	(Becker et al., 2006)	
Permethrin	53.65 (arithmetic)	659.15	129.1 (P90)	vacuum	Germany-Hannover	(Berger-Preiß et al., 2002)	
Phosmet	0.227	0.658		surface sampler	USA-Washington	(Simcox et al., 1995)	Non-farm
Phosmet	0.09	0.2		vacuum	USA-Washington	(Lu et al., 2000)	Non-farm
Phosmet	5.2	22		surface sampler	USA-Oregon	(Rothlein et al., 2006)	

Substance	Concer	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Pesticides							
						(Obendorf et al.,	
Picloram		1.2		vacuum	USA-New York state	2006)	
Resmethrin		0.8		vacuum	USA-New York state	(Obendorf <i>et al.</i> , 2006)	
Tetramethrin		0.4		vacuum	USA-New York state	(Obendorf <i>et al.</i> , 2006)	
trans-Chlordane <sup>2</sup>	5 μg/kg	0.073		airconditioning	Singapore	(Tan et al., 2007b)	
Trichloro-2-pyridinol <sup>4</sup>	0.535	0.950		surface sampler	USA-North Carolina	(Wilson <i>et al.</i> , 2003)	
Trifluralin		1.8		vacuum	USA-New York state	(Obendorf <i>et al.</i> , 2006)	
Phthalates							
Butylbenzyl phthalate (BBP)	157			vacuum	Sweden	(Bornehag <i>et al.</i> , 2004)	
Butylbenzyl phthalate (BBP)	86.1	815.7	218.5	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2004)	
Butylbenzyl phthalate (BBP)	319	45549	599	vacuum	Sweden	(Bornehag et al., 2005)	Dust from bedrooms
Butylbenzyl phthalate (BBP)	5.86	15.6		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Butylbenzyl phthalate (BBP)	84		416		Europe	(Wormuth <i>et al.</i> , 2006)	
Di(2-ethylhexyl) phthalate (DEHP)	508	5330	1840	vacuum	Germany	(Becker et al., 2004)	
Di(2-ethylhexyl) phthalate (DEHP)	775.5	1763	1542	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2004)	
Di(2-ethylhexyl) phthalate (DEHP)	741			vacuum	Sweden	(Bornehag et al., 2004)	
Di(2-ethylhexyl) phthalate (DEHP)	1310	40459	4069	vacuum	Sweden	(Bornehag et al., 2005)	

Substance	Concen	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric						
	mean	Maximum	P95				
Phthalates							
Di(2-ethylhexyl)							
phthalate (DEHP)	3214		7063	vacuum	Denmark	(Clausen et al., 2003)	Dust from schools
Di(2-ethylhexyl)							
phthalate (DEHP)			2600	vacuum	Germany	(Butte, 2001)	
Di(2-ethylhexyl)							
phthalate (DEHP)	858		2595	vacuum	Denmark	(Clausen et al., 2003)	
Di(2-ethylhexyl)							
phthalate (DEHP)			2000	vacuum	Germany	(Pohner et al., 1997)	
Di(2-ethylhexyl)							
phthalate (DEHP)	640			vacuum	Norway	(Oie et al., 1997)	
Di(2-ethylhexyl)						(Wormuth et al.,	
phthalate (DEHP)	1198		3470		Europe	2006)	
Diethyl phthalate							
(DEP)	44.6	632.2	159.6	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2004)	
Diethyl phthalate						(Wormuth et al.,	
(DEP)	26		114		Europe	2006)	
						(Wormuth et al.,	
Diisobutyl (DiBP)	84		130		Europe	2006)	
Diisodecyl (DIDP)	73		240		Europe	(Wormuth et al., 2006)	
					1	(Wormuth et al.,	
Diisononyl (DINP)	176		674		Europe	2006)	
Dimethyl phthalate	170		07.		Zurope	2000)	
(DMP)	10.8	157.9	46.4	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2004)	
Dimethyl phthalate					. , .	(Wormuth <i>et al.</i> ,	
(DMP)	1.1		1.8		Europe	2006)	
Dimethylpropyl					•	,	
phthalate (DMPP)	54.6	161.3	144.4	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2004)	
Di-n-butyl phthalate							
(DBP)	1.21	3.03		surface sampler	USA-North Carolina	(Wilson et al., 2003)	

Substance	Concer	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Phthalates							
Di-n-butyl phthalate (DBP)	178			vacuum	Sweden	(Bornehag <i>et al.</i> , 2004)	
Di-n-butyl phthalate (DBP)	55.6	141.4	129.6	vacuum	Germany-Berlin	(Fromme <i>et al.</i> , 2004)	
Di-n-butyl phthalate (DBP)	226	5446	568	vacuum	Sweden	(Bornehag et al., 2005)	
Di-n-butyl phthalate (DBP)	98		311		Europe	(Wormuth <i>et al.</i> , 2006)	
Brominated flame retardants (BFRs)	BFRs in μg/kg	BFRs in μg/kg					
BDE 100	1.15			vacuum	Kuwait	(Gevao et al., 2006)	
BDE 100	82,825			vacuum	Spain	(Regueiro <i>et al.</i> , 2006)	Average of 4 samples
BDE 100	274	2090		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 100	65	1200		airconditioning	Singapore	(Tan et al., 2007a)	
BDE 100	20.8	314		vacuum	Germany	(Knoth, 2003)	
BDE 100	490	21000	790	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 100	429				USA-Dallas	Schecter-2005	
BDE 138	17.3	111		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 138	37	2000	38	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 153	30.7	420		vacuum	Germany	(Knoth, 2003)	
BDE 153	1.31			vacuum	Kuwait	(Gevao et al., 2006)	
BDE 153	3,025			vacuum	Spain	(Regueiro et al., 2006)	Average of 4 samples
BDE 153	181	1510		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 154	13.8	210		vacuum	Germany	(Knoth, 2003)	

Substance	Concer	tration in dust	t (mg/kg)	Sampling	Location	Reference	Remarks
	Geometric						
	mean	Maximum	P95				
Brominated flame retardants (BFRs)							
BDE 154	1.39			vacuum	Kuwait	(Gevao et al., 2006)	
BDE 154	2,885			vacuum	Spain	(Regueiro <i>et al.</i> , 2006)	Average of 4 samples
BDE 154	156	1250		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 154	43	960		airconditioning	Singapore	(Tan et al., 2007a)	
BDE 154	380	18000	570	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 154	189				USA-Dallas	(Schecter et al., 2005)	
BDE 17	8.9	21.6		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 17	4.3	150	12	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 183	23.3	464		vacuum	Germany	(Knoth, 2003)	
BDE 183	3.57			vacuum	Kuwait	(Gevao et al., 2006)	
BDE 183	30.7	168		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 183	18	180		airconditioning	Singapore	(Tan et al., 2007a)	
BDE 183	44	650	200	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 183	19.3				USA-Dallas	(Schecter <i>et al.</i> , 2005)	
BDE 190	4.5	10.5		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 190	0.75	48	<loq< td=""><td>vacuum</td><td>Canada-Ottawa</td><td>(Wilford et al., 2005)</td><td></td></loq<>	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 196	14.5	38.6		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 197	17.4	77.2		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	

Substance	Concer	tration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Brominated flame retardants (BFRs)							
BDE 206	51.1	239		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 207	30	109		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 208	34.7	108		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 209	0.98	19100		vacuum	Germany	(Knoth, 2003)	
BDE 209	0.1288	0.338		vacuum	Kuwait	(Gevao et al., 2006)	
BDE 209	2.09	8750		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 209	2,200	13		airconditioning	Singapore	(Tan et al., 2007a)	
BDE 209	1.1	10	4.1	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 209	8,567				USA-Dallas	(Schecter et al., 2005)	
BDE 209	0.425				Europe	(Santillo et al., 2003) (Sjodin et al., 2004) (Al Bitar, 2004) (Knoth, 2003)	
BDE 209	4,028	65,777			USA	(Stapleton et al., 2005) (Schecter et al., 2005) (Sharp et al., 2004) (Costner et al., 2005)	
BDE 209	2,000	,			USA	(Sjodin <i>et al.</i> , 2004)	
BDE 209	0.732				Australia	(Sjodin <i>et al.</i> , 2006)	
BDE 209	0.619	2,230			Australia	(Toms et al., 2006)	
BDE 209	10,290	,			UK	(Santillo <i>et al.</i> , 2003) (Sjodin <i>et al.</i> , 2006)	
BDE 28	0.9	4.4		vacuum	Germany	(Knoth, 2003)	

Substance	Concer	ntration in dust	t (mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Brominated flame retardants (BFRs)							
BDE 28	0.35			vacuum	Kuwait	(Gevao et al., 2006)	
BDE 28	1.2	5.84		airconditioning	Singapore	(Tan et al., 2007a)	
BDE 28	15	550	28	vacuum	Canada-Ottawa	(Wilford <i>et al.</i> , 2005)	
BDE 28	20.3				USA-Dallas	(Schecter et al., 2005)	
BDE 33, 28	20.7	76.5		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 47	122	910		vacuum	Germany	(Knoth, 2003)	
BDE 47	27.2 ng/g				UK	(Santillo <i>et al.</i> , 2003) (Sjodin <i>et al.</i> , 2006)	
BDE 47	21.6 ng/g				Europe	(Santillo <i>et al.</i> , 2003) (Sjodin <i>et al.</i> , 2004) (Al Bitar, 2004; Knoth, 2003)	
BDE 47	6.57 ng/g	65 ng/g		vacuum	Kuwait	(Gevao et al., 2006)	
BDE 47	280			vacuum	Spain	(Regueiro <i>et al.</i> , 2006)	Average of 4 samples
BDE 47	1220	7610		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 47	110 ng/g	1500 ng/g		airconditioning	Singapore	(Tan et al., 2007a)	
BDE 47	55 ng/g	210 ng/g			Australia	(Toms et al., 2006)	
BDE 47	60 ng/g				Australia	(Sjodin et al., 2006)	
BDE 47	1100 ng/g	33000 ng/g	2600	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
						(Sjodin et al., 2004)	
BDE 47	230 ng/g				USA		

Substance	Concer	Concentration in dust (mg/kg)			Location	Reference	Remarks	
	Geometric mean	Maximum	P95	Sampling				
Brominated flame retardants (BFRs)								
BDE 47	1595 ng/g	10538 ng/g			USA	(Stapleton <i>et al.</i> , 2005)(Schecter <i>et al.</i> , 2005) (Sharp <i>et al.</i> , 2004) (Costner <i>et al.</i> , 2005)		
BDE 47	1621	10336 lig/g			USA-Dallas	(Schecter <i>et al.</i> , 2005)		
BDE 49	25.8	282		vacuum	Germany	(Knoth, 2003)		
BDE 66	28.5	142		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)		
BDE 66	37	1800	45	vacuum	Canada-Ottawa	(Wilford et al., 2005)		
BDE 85	7.1	74.7		vacuum	Germany	(Knoth, 2003)		
BDE 85	0.72			vacuum	Kuwait	(Gevao et al., 2006)		
BDE 85	1,365			vacuum	Spain	(Regueiro <i>et al.</i> , 2006)	Average of 4 samples	
BDE 85	83.4	620		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)		
BDE 85	190	9700	200	vacuum	Canada-Ottawa	(Wilford et al., 2005)		
BDE 99	180 ng/g	2850		vacuum	Germany	(Knoth, 2003)		
BDE 99	79.8 ng/g				UK	(Santillo <i>et al.</i> , 2003) (Sjodin <i>et al.</i> , 2006)		
BDE 99	32.8 ng/g				Europe	(Santillo et al., 2003) (Sjodin et al., 2004) (Al Bitar, 2004) (Knoth, 2003)		
BDE 99	1977 ng/g	13841 ng/g			USA	(Stapleton <i>et al.</i> , 2005; Schecter <i>et al.</i> , 2005) (Sharp <i>et al.</i> , 2004) (Costner <i>et al.</i> , 2005)		

Substance	Concer	ntration in dust	(mg/kg)	Sampling	Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Brominated flame retardants (BFRs)							
BDE 99	6.00 ng/g	36 ng/g		vacuum	Kuwait	(Gevao et al., 2006)	
BDE 99	79.8 ng/g	294 ng/g			Australia	(Toms et al., 2006)	
BDE 99	880 ng/g				USA	(Sjodin et al., 2004)	
BDE 99	106 ng/g				Australia	(Sjodin et al., 2006)	
BDE 99	31,525			vacuum	Spain	(Regueiro et al., 2006)	Average of 4 samples
BDE 99	1700	13800		vacuum	USA-Washington	(Stapleton <i>et al.</i> , 2005)	
BDE 99	340 ng/g	6300 ng/g		airconditioning	Singapore	(Tan et al., 2007a)	
BDE 99	1800 ng/g	60000 ng/g	4700	vacuum	Canada-Ottawa	(Wilford et al., 2005)	
BDE 99	2295				USA-Dallas	(Schecter <i>et al.</i> , 2005)	
BDE99	8.5 ng/g	7771			Netherlands	(De Boer, 2007)	
BDE153	76	1400		airconditioning	Singapore	(Tan et al., 2007a)	
BDE153	470	25000	520	vacuum	Canada-Ottawa	(Wilford <i>et al.</i> , 2005)	
BDE153	199				USA-Dallas	(Schecter <i>et al.</i> , 2005)	
Polycyclic aromatic h	nydrocarbons (PA	Hs)					
Acenaphthene	0.05	0.18		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
Acenaphthene	0.032	1.9		different methods		(Maertens <i>et al.</i> , 2004)	Average of 18 studies
Acenaphthene	0.008	0.019		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Acenaphthylene	0.08	0.27		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
Acenaphthylene	0.026	0.52		different methods		(Maertens <i>et al.</i> , 2004)	Average of 18 studies
Acenaphthylene	0.006	0.023		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Anthracene	0.12	0.75		surface sampler	USA-North Carolina	(Chuang et al., 1999)	

Substance	Concentration in dust (mg/kg)			Sampling	Location	Reference	Remarks
	Geometric			1			
	mean	Maximum	P95				
Polycyclic aromatic							
hydrocarbons (PAHs)							
				different		(Maertens et al.,	
Anthracene	0.065	5.8		methods		2004)	Average of 18 studies
Anthracene	0.017	0.066		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Benz[a]anthracene	0.22	0.69		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
				different		(Maertens et al.,	
Benz[a]anthracene	0.241	40		methods		2004)	Average of 18 studies
Benz[a]anthracene	0.090	0.519		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Benzo[a]pyrene	0.23	0.63		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
				different		(Maertens et al.,	
Benzo[a]pyrene	0.285	54	13	methods		2004)	Average of 18 studies
Benzo[a]pyrene	0.134	0.768		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Benzo[b,k]fluoranthene	0.55	1.34		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
				different		(Maertens et al.,	
Benzo[b,k]fluoranthene	0.570	108		methods		2004)	Average of 18 studies
Benzo[b,k]fluoranthene	0.253	1.44		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Benzo[b,k]fluoranthene	0.050	0.496		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Benzo[e]pyrene	0.26	0.75		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
Benzo[e]pyrene	0.286	41		different methods		(Maertens et al., 2004)	Average of 18 studies
Benzo[e]pyrene	0.144	0.809		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Benzo[g,h,i]perylene	0.25	0.61		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
10/ / 11 3				different		(Maertens et al.,	
Benzo[g,h,i]perylene	0.252	35		methods		2004)	Average of 18 studies
Benzo[g,h,i]perylene	0.173	0.961		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Biphenyl	0.002	0.005		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Chrysene	0.39	2.41		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
				different		(Maertens <i>et al.</i> ,	
Chrysene	0.372	43		methods		2004)	Average of 18 studies
Chrysene	0.169	0.838		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Coronene	0.13	0.50		surface sampler	USA-North Carolina	(Chuang et al., 1999)	

Concentration in dust (mg/kg)			Sampling	Location	Reference	Remarks
Geometric						
mean	Maximum	P95				
			different		(Maertens et al.,	
0.095	7.2		methods		2004)	Average of 18 studies
0.067	0.32		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
0.08	0.22		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
			different		(Maertens et al.,	
0.034	0.62		methods		2004)	Average of 18 studies
0.035	0.172		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
0.10	0.41		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
			different		(Maertens et al.,	
0.082	9.0		methods		2004)	Average of 18 studies
0.060	0.294		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
0.52	1.89		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
			different		(Maertens et al.,	
0.588	90		methods		2004)	Average of 18 studies
0.297	1.56		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
0.12	1.22		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
			different		(Maertens et al.,	
0.054	3.0		methods		2004)	Average of 18 studies
0.011	0.028		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
0.23	0.70		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
			different		(Maertens et al.,	
0.255	41		methods		2004)	Average of 18 studies
•			*			
0.33	4.30		surface sampler	USA-North Carolina	(Chuang <i>et al.</i> , 1999)	
	0.095 0.067 0.08 0.034 0.035 0.10 0.082 0.060 0.52 0.588 0.297 0.12 0.054 0.011 0.23 0.255 0.169	Geometric mean         Maximum           0.095         7.2           0.067         0.32           0.08         0.22           0.034         0.62           0.035         0.172           0.10         0.41           0.082         9.0           0.060         0.294           0.52         1.89           0.588         90           0.297         1.56           0.12         1.22           0.054         3.0           0.011         0.028           0.23         0.70           0.255         41           0.169         0.963	Geometric mean         Maximum         P95           0.095         7.2           0.067         0.32           0.08         0.22           0.034         0.62           0.035         0.172           0.10         0.41           0.082         9.0           0.060         0.294           0.52         1.89           0.588         90           0.297         1.56           0.12         1.22           0.054         3.0           0.011         0.028           0.23         0.70           0.255         41           0.169         0.963	Geometric mean         Maximum         P95           0.095         7.2         different methods           0.067         0.32         surface sampler           0.08         0.22         surface sampler different methods           0.034         0.62         methods           0.035         0.172         surface sampler different methods           0.082         9.0         methods           0.060         0.294         surface sampler different methods           0.52         1.89         surface sampler different methods           0.297         1.56         surface sampler different methods           0.012         1.22         surface sampler different methods           0.011         0.028         surface sampler different methods           0.23         0.70         surface sampler different methods           0.255         41         methods	Geometric mean         Maximum         P95           0.095         7.2         different methods           0.067         0.32         surface sampler USA-North Carolina           0.08         0.22         surface sampler USA-North Carolina           0.034         0.62         methods           0.035         0.172         surface sampler USA-North Carolina           0.10         0.41         surface sampler different methods           0.082         9.0         methods           0.060         0.294         surface sampler USA-North Carolina           0.52         1.89         surface sampler USA-North Carolina           0.588         90         different methods           0.297         1.56         surface sampler USA-North Carolina           0.12         1.22         surface sampler USA-North Carolina           0.054         3.0         methods           0.011         0.028         surface sampler USA-North Carolina           0.23         0.70         surface sampler USA-North Carolina           0.255         41         methods           0.169         0.963         surface sampler USA-North Carolina	Description

Substance	Concen	Concentration in dust (mg/kg)			Location	Reference	Remarks
	Geometric mean	Maximum	P95				
Polycyclic aromatic hydrocarbons (PAHs)							
Naphthalene	0.068	42		different methods		(Maertens et al., 2004)	Average of 18 studies
Naphthalene	0.010	0.035		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Phenanthrene	0.44	2.15		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
Phenanthrene	0.416	43		different methods		(Maertens <i>et al.</i> , 2004)	Average of 18 studies
Phenanthrene	0.144	0.596		surface sampler	USA-North Carolina	(Wilson et al., 2003)	
Pyrene	0.43	1.65		surface sampler	USA-North Carolina	(Chuang et al., 1999)	
Pyrene	0.490	69		different methods		(Maertens et al., 2004)	Average of 18 studies
Pyrene	0.229	1.20		surface sampler	USA-North Carolina	(Wilson et al., 2003)	

- Appendix 1 Remarks

  1. Alpha-Chlordane = cis-Chlordane.
- 2. Gamma-Chlordane = trans-Chlordane.
- Lindane is technical gamma-HCH.
   Trichloro-2-pyridinol is a metabolite of chlorpyrifos.



### Appendix 2 List of TDIs and background exposure estimates

Table Appendix 2. TDIs and background exposures

	TDI	Reference	Backş	ground	Specification background	Reference
	(μg/kg bw/day)		(μg/kg	bw/day)		
			Adult	Child		
Metals						
Aluminium	750	(Van Engelen <i>et al.</i> , in press)	80-180	300	food, drinking- water	(Van Engelen <i>et al.</i> , in press)
Antimony	6	(Van Engelen <i>et al.</i> , in press)	0.018- 0.48	0.53	food, drinking- water	(Van Engelen <i>et al.</i> , in press)
Arsenic	1	(Van Engelen <i>et al.</i> , in press)	0.3	0.4-0.7	food, drinking- water	(Van Engelen <i>et al.</i> , in press) (Baars <i>et al.</i> , 2001)
Barium	600	(Van Engelen <i>et al.</i> , in press)		9	food, drinking- water	(Van Engelen <i>et al.</i> , in press)
Beryllium	0.5	(Janssen <i>et al.</i> , 1995)	C	0.3	food, drinking- water	(Janssen <i>et al.</i> , 1995)
Bismuth	n.a.		unk	nown		
Cadmium	0.5	(Oomen et al., 2007)	0.	.45	food, drinking- water	(Van Engelen <i>et al.</i> , in press)
Chromium III soluble	5.0			1	food, drinking- water	(Baars et al., 2001)
Chromium III insoluble	5000	(Baars et al., 2001)		1	food, drinking- water	(Baars et al., 2001)
Chromium VI	5	(Baars et al., 2001)	5.7×10 <sup>-6</sup>	-0.43×10 <sup>-3</sup>	air	(Baars et al., 2001)
Cobalt	1.4	(Van Engelen <i>et al.</i> , in press)		0.6	food, drinking- water	(Van Engelen <i>et al.</i> , in press)

					Specification		
	TDI	Reference	Background		background	Reference	
	(μg/kg bw/day)		(μg/kg	bw/day)			
			Adult	Child			
Metals							
					food, drinking-		
Copper	83	(Van Engelen et al., in press)		60	water	(Van Engelen et al., in press)	
Lead	3.6	(Van Engelen <i>et al.</i> , in press)	1.11	1.83		(Gezondheidsraad (Dutch Health Council), 1997)	
					food, drinking-		
Lithium	n.a.		unk	nown	water		
					food, drinking-		
Magnesium	6700	(Food Standards Agency)	4	600	water	(Food Standards Agency)	
					food, drinking-		
Manganese	160	(Van Engelen <i>et al.</i> , in press)	1	130	water	(Van Engelen <i>et al.</i> , in press)	
					food, dental		
Mercury	2	(Van Engelen <i>et al.</i> , in press)	(	0.1	fillings	(Van Engelen <i>et al.</i> , in press)	
					food, drinking-		
Molybdenum	10	(Baars et al., 2001)		4	water	(Baars et al., 2001)	
	10				food, drinking-	(Van Engelen <i>et al.</i> , in press)	
Nickel	10	(Van Engelen <i>et al.</i> , in press)	4	8	water	(Baars et al., 2001)	
Rubidium	n.a.		unk	nown			
					food, drinking-		
Selenium	5	(Van Engelen <i>et al.</i> , in press)		2	water	(Van Engelen <i>et al.</i> , in press)	
					food, drinking-		
Silver	5	(Van Engelen <i>et al.</i> , in press)		1.3	water	(Van Engelen et al., in press)	
					food, drinking-		
Strontium	600	(Van Engelen et al., in press)		18	water	(Van Engelen et al., in press)	
Tellurium	2	(Janssen et al., 1998)		1.4	food	(Janssen <i>et al.</i> , 1998)	
Thallium	0.2	(Janssen et al., 1998)	0	0.03	food	(Janssen et al., 1998)	
Tin	2000	(Van Engelen <i>et al.</i> , in press)	2	290	food, water	(Van Engelen <i>et al.</i> , in press)	

				Specification	
	TDI	Reference	Background	background	Reference
	(μg/kg bw/day)		(μg/kg bw/day)		
			Adult Child		
Metals		T		T	
Titanium	12000	(Janssen and Baars, 2004)	7	food	(Janssen and Baars, 2004)
Tungsten	n.a.		unknown		
				food, drinking-	
Uranium	2	(ATSDR, 1999)	0.06	water	(Bruggen et al., 1998)
Vanadium	2		0.3	food	(Janssen et al., 1998)
Zinc	500	(Van Engelen et al., in press)	350	food	(Van Engelen et al., in press)
Organotin compounds					
Dibutyltin (DBT)					
Dioctyltin (DOT)					
Monobutyltin (MBT)					
Monooctyltin (MOT)					
Tributyltin (TBT)					
				food, drinking-	
Sum organotins	0.25	(Van Engelen <i>et al.</i> , in press)	0.083	water	(Van Engelen <i>et al.</i> , in press)
Pesticides		I			1
2.4-D	10	(IMPR 2001)	unknown	food, drinking-	
		(JMPR, 2001)		water	
Alachlor	10	(USEPA, 1998)	unknown	food	

	TDI	Reference	Background	Specification background	Reference
		Kererence	•	Dackground	Reference
	(μg/kg bw/day)		(μg/kg bw/day)		
			Adult Child		
Pesticides					
Aldrin	0.1 (som aldrin+dieldrin)	(Baars <i>et al.</i> , 2001)	< 0.04	food	(Baars et al., 2001)
alpha-Chlordane	0.5	(Janssen et al., 1995)	unknown	food	
alpha-HCH	1	(Baars et al., 2001)	< 0.03	food	(Baars et al., 2001)
Atrazine	35	(USEPA, 1993)	unknown	food	
				food, drinking-	
Azinphos methyl	5	(Janssen et al., 1995)	unknown	water	
				food, drinking-	
beta-HCH	0.02	(Baars et al., 2001)	< 0.01	water	(Baars et al., 2001)
				food, drinking-	
Carbaryl	8	(JMPR, 2001)	unknown	water	
				food, drinking-	
Chloroprofam	50	(JMPR, 2005)	unknown	water	
				food, drinking-	
Chlorpyrifos	10	(JMPR, 2004a)	unknown	water	
	0.5 (som DDT,			food, drinking-	
DDD	DDD, DDE)	(Baars et al., 2001)	unknown	water	
	0.5 (som DDT,			food, drinking-	
DDE	DDD, DDE)	(Baars et al., 2001)	unknown	water	
	0.5 (som DDT,			food, drinking-	
DDT	DDD, DDE)	(Baars et al., 2001)	unknown	water	
				food, drinking-	
delta-HCH	n.a.		unknown	water	
				food, drinking-	
Diazinon	5	(JMPR, 2006)	unknown	water	

	TDI	TDI Reference		Specification background	Reference	
		Keierence	Background	Dackground	Reference	
	(μg/kg bw/day)		(μg/kg bw/day)			
			Adult Child			
Pesticides						
				food, drinking-		
Dicamba	125	(Hoeven and Engelen, 1997)	unknown	water		
	0.1 (som					
Dieldrin	aldrin+dieldrin)	(Baars et al., 2001)	< 0.04	food	(Baars et al., 2001)	
Dimethyl						
organiphosphate						
				food, drinking-		
Endrin	0.2	(Baars et al., 2001)	< 0.04	water	(Baars et al., 2001)	
				food, drinking-		
Ethyl parathion	4	(JMPR, 1995)	unknown	water		
				food, drinking-		
Flutolanil	90	(JMPR, 2002)	unknown	water		
				food, drinking-		
gamma-Chlordane	0.5	(Janssen et al., 1995)	unknown	water		
				food, drinking-		
Glyphosate	1000	(JMPR, 2004b)	unknown	water		
				food, drinking-		
Heptachlor	0.1	(Janssen et al., 1995)	0.001	water	(Janssen et al., 1995)	
				food, drinking-		
Lindane [= gamma-HCH]	0.04	(Baars et al., 2001)	< 0.03	water	(Baars et al., 2001)	
				food, drinking-		
Malathion	300	(JMPR, 2003)	unknown	water		
				food, drinking-		
Mecoprop	3.33	(WHO, 1996)	unknown	water		

	TDI	Reference	Background	Specification background	Reference
	(μg/kg bw/day)		(μg/kg bw/day		
	(μg/kg bw/day)		Adult Child		
Pesticides			Addit Clind		
				food, drinking-	
Methamidophos	4	(JMPR, 2002)	unknown	water	
<b>'</b>				food, drinking-	
Methyl parathion	3	(JMPR, 1995)	unknown	water	
<del> </del>				food, drinking-	
Metolachlor	3.5	(WHO, 2003)	unknown	water	
				food, drinking-	
Pendimethalin	125	(EU, 2003)	unknown	water	
				food, drinking-	
Permethrin	50	(JMPR, 2002)	unknown	water	
				food, drinking-	
Phosmet	3	(EU, 2006)	unknown	water	
				food, drinking-	
Picloram	200	(BfR, 2006)	unknown	water	
				food, drinking-	
Resmethrin	30	(JMPR, 1991)	unknown	water	
				food, drinking-	
Tetramethrin	20	(AG, 1992)	unknown	water	
				food, drinking-	
Trichloro-2-pyridinol	n.a.		unknown	water	
				food, drinking-	
Trifluralin	15	(BfR, 2006)	unknown	water	
				food, drinking-	
	10	(JMPR, 1995)	unknown	water	

	TDI		D. d.		Specification	Reference
	TDI	Reference		ground	background	Reference
	(μg/kg bw/day)		(μg/kg bw/day)			
			Adult	Child		
Phthalates						
Butylbenzyl phthalate					food, drinking-	
(BBP)	500	(EFSA, 2005a)	5.0-9.0		water	(Baars et al., 2001)
Di(2-ethylhexyl)phthalate					food, drinking-	
(DEHP)	50	(EFSA, 2005b)	3-16	12-26	water	(EFSA, 2005b)
					food, drinking-	
Diethyl phthalate (DEP)	200	(Baars et al., 2001)	unk	nown	water	
					food, drinking-	
Diisobutyl (DiBP)	n.a.		unk	nown	water	
					food, drinking-	
Diisodecyl (DIDP)	150	(EFSA, 2005c)		7	water	(EFSA, 2005c)
					food, drinking-	
Diisononyl (DINP)	150	(EFSA, 2005d)		10	water	(EFSA, 2005d)
Dimethyl phthalate					food, drinking-	
(DMP)	n.a.		unk	nown	water	
Dimethylpropyl phthalate					food, drinking-	
(DMPP)	n.a.		unk	nown	water	
Di-n-butyl phthalate					food, drinking-	
(DBP)	52	(Baars et al., 2001)	unk	nown	water	
Brominated flame retardar	nts (BFRs)					
BDE 47						
BDE 99	0.00026	(Winter-Sorkina et al., 2006)				
BDE 100						

					Specification	
	TDI	Reference	Back	ground	background	Reference
	(μg/kg bw/day)		(μg/kg	bw/day)		
			Adult	Child		
Brominated flame retarda	nts (BFRs)	,				
BDE 183						
BDE 209						
Polycyclic aromatic hydro	carbons (PAHs)					
Acenaphthene					food	
Acenaphthylene					food	
Anthracene	40	(Baars et al., 2001)			food	
Benz[a]anthracene					food	
	0.0005 (BaP-		in EU vi			
Benzo[a]pyrene	indicator approach)	(Baars et al., 2001) (SCF 2002)	0.0006 (	BaP)	food	(SCF, 2002)
Benzo[b,k]fluoranthene					food	
Benzo[e]pyrene					food	
Benzo[g,h,i]perylene	30	(Baars et al., 2001)			food	
Biphenyl	50	(USEPA, 1989)			food	
Chrysene					food	
Coronene					food	
Cyclopenta[c,d]pyrene					food	
Dibenzo[a,h]anthracene					food	
Fluoranthene					food	
Fluorene	40	(Baars et al., 2001)			food	
Indeno[1,2,3-c,d]pyrene					food	
Naphthalene	40	(Baars et al., 2001)			food	

	TDI	Reference	Background		Specification background	Reference			
	(μg/kg bw/day)		(μg/kg bw/day)						
			Adult	Child					
Polycyclic aromatic hydrocarbons (PAHs)									
Phenanthrene	40	(Baars et al., 2001)			food				
Pyrene					food				

# RIVM National Institute for Public Health and the Environment P.O. Box 1 3720 BA Bilthoven

The Netherlands www.rivm.com